



# Hydroxyl radicals from secondary organic aerosol decomposition in water

Haijie Tong<sup>1</sup>, Andrea M. Arangio<sup>1</sup>, Pascale S. J. Lakey<sup>1</sup>, Thomas Berkemeier<sup>1</sup>, Fobang Liu<sup>1</sup>, Christopher J. Kampf<sup>1,2</sup>, William H. Brune<sup>3</sup>, Ulrich Pöschl<sup>1</sup>, and Manabu Shiraiwa<sup>1</sup>

<sup>1</sup>Multiphase Chemistry Department, Max Planck Institute for Chemistry, Mainz, Germany

<sup>2</sup>Institute for Inorganic and Analytical Chemistry, Johannes Gutenberg University Mainz, Mainz, Germany

<sup>3</sup>Department of Meteorology, Pennsylvania State University, University Park, PA 16802, USA

Correspondence to: Manabu Shiraiwa (m.shiraiwa@mpic.de)

Received: 22 October 2015 – Published in Atmos. Chem. Phys. Discuss.: 3 November 2015

Revised: 29 January 2016 – Accepted: 2 February 2016 – Published: 15 February 2016

**Abstract.** We found that ambient and laboratory-generated secondary organic aerosols (SOA) form substantial amounts of OH radicals upon interaction with liquid water, which can be explained by the decomposition of organic hydroperoxides. The molar OH yield from SOA formed by ozonolysis of terpenes ( $\alpha$ -pinene,  $\beta$ -pinene, limonene) is  $\sim 0.1\%$  upon extraction with pure water and increases to  $\sim 1.5\%$  in the presence of  $\text{Fe}^{2+}$  ions due to Fenton-like reactions. Upon extraction of SOA samples from OH photooxidation of isoprene, we also detected OH yields of around  $\sim 0.1\%$ , which increases upon addition of  $\text{Fe}^{2+}$ . Our findings imply that the chemical reactivity and aging of SOA particles is strongly enhanced upon interaction with water and iron. In cloud droplets under dark conditions, SOA decomposition can compete with the classical  $\text{H}_2\text{O}_2$  Fenton reaction as the source of OH radicals. Also in the human respiratory tract, the inhalation and deposition of SOA particles may lead to a substantial release of OH radicals, which may contribute to oxidative stress and play an important role in the adverse health effects of atmospheric aerosols.


## 1 Introduction

Secondary organic aerosols (SOA) account for a major fraction of fine air particulate matter and have a strong influence on climate and public health (Jimenez et al., 2009; Pöschl et al., 2010; Huang et al., 2014). Formation of SOA is triggered by oxidation of volatile organic compounds followed by condensation of semi-volatile oxidation products (Hallquist et

al., 2009; Donahue et al., 2012). Recently, it has been shown that extremely low volatility organic compounds contribute significantly to SOA growth (Ehn et al., 2014; Jokinen et al., 2015; Mentel et al., 2015).

Particle phase chemistry and cloud processing are also efficient pathways for SOA formation and aging (Kalberer et al., 2004; Herrmann et al., 2005; Ervens et al., 2011; Shiraiwa et al., 2013). Evolution of SOA is one of the largest uncertainties in the current understanding of air quality, climate and public health (Kanakidou et al., 2005; Solomon, 2007). With regard to SOA health effects, substantial amounts of reactive oxygen species including organic radicals are detected in ambient and laboratory-generated SOA (Venkatachari and Hopke, 2008; Chen and Hopke, 2010; Chen et al., 2010; Fuller et al., 2014). Despite intensive research, multiphase chemical reactions of SOA in the atmosphere and upon interaction with the human respiratory tract are not well understood (Pöschl and Shiraiwa, 2015).

OH radicals in atmospheric droplets originate from the uptake of gaseous OH radicals (Jacob, 1986; Arakaki et al., 2013) as well as photolysis of ozone (Anglada et al., 2014). A recent study has shown that SOA can form OH radicals in the aqueous phase under light conditions (Badali et al., 2015). Under dark conditions, Fenton reactions between  $\text{H}_2\text{O}_2$  and iron ions have been regarded as the main source of OH radicals so far (Herrmann et al., 2005). In this study, we found that OH radicals are formed by decomposition of SOA upon interactions of water and iron ions under dark conditions.



**Figure 1.** Schematics of the experimental setup for generation and collection of SOA particles.

## 2 Methods

### 2.1 SOA formation and particle collection

Figure 1 shows the experimental setup for generation of secondary organic aerosols (SOA).  $O_3$  was used as oxidant for oxidation of  $\alpha$ -pinene,  $\beta$ -pinene and limonene, and OH radicals were used for naphthalene.  $O_3$  was generated via synthetic air (Westfalen AG,  $1.8\text{--}2.1\text{ L min}^{-1}$ ) passing through a 185 nm UV light ( $O_3$  generator, L.O.T.-Oriol GmbH & Co. KG). The typical ozone concentrations were 600 ppb for  $\alpha$ -pinene,  $\beta$ -pinene and limonene, and 1200 ppb for naphthalene. A total of 1 mL of  $\alpha$ -pinene (98 %, Sigma Aldrich),  $\beta$ -pinene (99 %, Sigma Aldrich) or limonene (99 %, Sigma Aldrich) was kept in a 1.5 mL amber glass vial (VWR International GmbH), and 5–10 g of naphthalene crystals (99.6 %, Alfa Aesar GmbH & Co. KG) was put in a 100 mL glass bottle (DURAN Group GmbH) as SOA precursor sources. A total of 1 bar and  $50\text{--}150\text{ ccm min}^{-1}$   $N_2$  (99.999 %, Westfalen AG) flow was passed through these sources, and the evaporated volatile organic compound (VOC) vapours were introduced into a 7 L quartz flow tube reactor for gas-phase oxidation reaction with  $O_3$  or OH radicals with a reaction time of  $\sim 3$  min. SOA by  $\alpha$ -pinene,  $\beta$ -pinene and limonene were generated under dark and dry conditions. The flow tube reactor is surrounded by four UV lights (wavelength of 254 nm, LightTech Lamp Technology Ltd.), which were turned on to generate OH radicals by photolysis of ozone and water vapour. The relative humidity in the flow tube was 30 % for generating naphthalene SOA, and other experiments were conducted under dry conditions. Isoprene SOA was produced in a potential aerosol mass (PAM) chamber through the reaction of gas phase OH radicals and isoprene. The detailed information about this chamber has been described elsewhere (Kang et al., 2007; Lambe et al., 2011), and the SOA generated by the PAM chamber have been shown to be similar to SOA generated in large environmental chambers (Bruns et al., 2015; Lambe et al., 2015) and the atmosphere (Ortega et al., 2015) in terms of oxidation state and chemical composition. Briefly the isoprene vapour was taken into the chamber by  $N_2$  gas with an estimated concentration of tens of parts per million (ppm). Ozone concentration in the PAM was 6–15 ppm, and relative humidity was 30–40 %.

Number concentration and size distribution of the generated SOA particles were characterized using a scanning mobility particle sizer (SMPS, GRIMM Aerosol Technik GmbH & Co. KG). The typical size of the SOA ranged from 50 to 400 nm. The median diameters of the mass size distribution were 100–200 nm.  $MnO_2$  (copper mesh covered with  $MnO_2$  from ANSYCO Analytische Systeme und Komponenten GmbH fixed in Gelman filter) and charcoal (4–8 mesh, Sigma Aldrich) denuders were used to remove unreacted  $O_3$  before the collection of SOA particles on a filter. SOA was collected on 47 mm Omnipore Teflon filters (100 nm pore size, Merck Chemicals GmbH). The concentration of  $O_3$  was monitored after an ozone denuder with an ozone analyser (typically 0–20 ppb, model 49i, Thermo Fisher Scientific Inc.). Two silica gel (2–4 mm, Carl Roth GmbH & Co. KG) denuders were used to dry the naphthalene SOA before collection.

Blank tests confirmed that no radicals were produced without SOA particles on a filter. Condensation of water vapour on a filter during SOA collection was negligible. A Teflon filter with particle loading was weighed using a XSE105DU balance with accuracy of  $\pm 20\text{ }\mu\text{g}$ . It was then immersed into a 0.5–1 mL 10 mM BMPO water solution and stirred with a vortex shaker (Heidolph Reax 1) for 2–7 min for particle extraction. A typical extraction efficiency of  $> 70\%$  in weight can be obtained with 7 min extraction time. After extraction, the filter was dried under 2–3 bar  $N_2$  for  $\sim 10$  min and the filter was weighed. The weight difference was regarded as the weight of extracted particles. The final SOA concentration depends on the extraction time, and the average molar mass of SOA was assumed to be  $200\text{ g mol}^{-1}$  in calculating SOA concentrations. The pH of SOA solutions was in the range of 4.8–6.4.

A micro-orifice uniform deposit impactor (MOUDI, 110-R mode, MSP Corporation) was used for collection of ambient particles on the roof of the Max Planck Institute for Chemistry (Mainz, Germany) in 24 h time resolution with a flow rate of  $30\text{ L min}^{-1}$  from 17:30 UTC + 1 4 June 2015 to 17:30 5 June 2015 and from 17:30 7 June 2015 to 17:30 8 June 2015. Particles within the diameter range of 180–320 nm, which is the size range dominated by organic aerosols in Mainz (Faber et al., 2013), were used for further analysis. The mass loading of these two samples on filters were  $\sim 70$  and  $80\text{ }\mu\text{g}$ , respectively. Teflon filters of 47 mm diameter (100 nm pore size, Merck Chemicals GmbH) were


used to collect the roof particles. Filters were cleaned with pure ethanol and ultra-pure water and dried by nitrogen gas before sampling and weighing. The extraction procedure is the same as that for laboratory SOA, and the field particle extracts were concentrated with a  $N_2$  flux to obtain high signal-to-noise ratio spectra. Concentrations of field particles in water extracts for EPR measurements were  $\sim 0.3 \text{ g L}^{-1}$ , which is of the same order of magnitude as extracts of laboratory-generated SOA.

## 2.2 CW-EPR

Continuous-wave electron paramagnetic resonance (CW-EPR) spectroscopy (EMXplus-10/12, Bruker, Germany) was applied for detection of radicals. A total of 15–30  $\mu\text{L}$  sample solutions were kept in a 50  $\mu\text{L}$  capacity micropipette and inserted into a highly sensitive cavity (E4119001 HS-W1) for analysis. The set of EPR parameters used for this study was as follows: a modulation frequency of 100 kHz; a modulation amplitude of 0.6 or 1; microwave power of 2.149 mW (20 dB) or 21.17 mW (10 dB); a receiver gain of 40 dB; a time constant of 0.01 ms; and a magnetic field scan of 100 G. After the SOA extraction, the samples were immediately analysed by an EPR.

The spin trap 5-*tert*-butoxycarbonyl-5-methyl-1-pyrroline-N-oxide (BMPO, high purity, Enzo Life Sciences GmbH) was used as a trapping agent of OH radicals. Compared to other spin-trapping agents such as 5, 5-dimethyl-1-pyrroline N-oxide (DMPO), BMPO has the following advantages: high purity and stability in the crystalline phase; highly distinguishable EPR spectra for different structure of the trapped radicals; and spectra with high signal-to-noise ratio. Buffer solutions are often used in the spin-trapping technique, but they were not used in this study to avoid changing the real acidity environment of SOA solutions. A BMPO concentration of 10 mM was used. No significant difference was observed among 10, 20, 30, 40 and 50 mM BMPO solutions, confirming that a BMPO concentration of 10 mM is sufficient to achieve the maximum trapping efficiency. The influence of the BMPO concentration on the aqueous phase OH radical trapping efficiency for  $\beta$ -pinene SOA was investigated as shown in Fig. S3. Further blank tests confirmed that  $H_2O_2$  (30 %, Sigma Aldrich),  $Fe^{2+}$  and  $Fe^{3+}$  ( $Fe_2O_3 \cdot xH_2O$ , 97 %, Sigma Aldrich) do not induce OH radical formation when each of them is mixed with BMPO in water (Fig. S4).

The spin-counting method was applied for quantification of OH radicals using the embedded subroutine of the Bruker Xenon software (Weber, 2012). For better quantification of detected radicals, the spin-fitting method (Bruker Xenon software, chapter 13; Weber, 2012) was used to increase the signal-to-noise ratio especially for low radical concentrations. The required parameters are hyperfine splitting parameters for OH radicals, which were taken from Zhao et al. (2001). Spectral simulations for radical adducts were




**Figure 2.** EPR spectra of sample solutions mixed with the spin-trapping agent BMPO: (a)  $\alpha$ -pinene SOA, (b)  $\beta$ -pinene SOA, (c) limonene SOA, (d) isoprene SOA, (e) naphthalene SOA, (f) 180–320 nm size field particles, (g) 180–320 nm size field particles mixed with  $Fe^{2+}$ , (h) *tert*-butyl hydroperoxide solution and (i)  $H_2O_2$  solution with  $Fe^{2+}$ . The four peaks (dotted lines) are characteristic of BMPO-OH adducts.

carried out using the Matlab-based computational package Easyspin (Stoll and Schweiger, 2006). A global optimization (genetic algorithm) was conducted to obtain parameters for simulating the EPR spectrum. The parameter set was further optimized using the particle swarm method within the Easyspin program. The function “garlic” for cw EPR spectra in isotropic and fast motion regimes was chosen for simulation. The hyperfine splitting constants for simulation were taken from the Zhu et al. (2009).

## 2.3 LC-MS/MS

The SOA extracts mixed with spin-trapping agent BMPO were also analysed with a nanoHPLC-chip-MS/MS system (Agilent), which consists of a nano pump (G2226A) with four-channel micro-vacuum degasser (G1379B), a microfluidic chip cube with electrospray ionization (ESI) source (G4240-62010) interfaced to a Q-TOF mass spectrometer (6540; nominal mass resolution 30 000 at a scan rate of  $5 \text{ s}^{-1}$ ), a capillary pump (G1376A) with degasser (G1379B), and an auto-sampler with thermostat (G1377A). All modules were controlled by Mass Hunter software (Rev. B.05.01, Agilent). Eluents used were 3 % (*v/v*) acetone nitrile (Chromasolv, Sigma, Seelze, Germany) in water/formic acid (0.1 % *v/v*, Chromasolv, Sigma, Seelze, Germany) (Eluent A) and 3 % water/formic acid (0.1 % *v/v*) in acetone nitrile (Eluent B). The flow rate was  $400 \text{ nL min}^{-1}$  with a gradient program that starting with 3 % B for 3 min followed by a 36 min step that raised eluent B to 60 %. Further, eluent B was increased to 80 % at 40 min and returned to initial condi-



**Figure 3.** LC-MS/MS analysis. (a) LC-MS chromatogram of aqueous BMPO solution (black line) and BMPO mixed with  $\beta$ -pinene SOA water extracts (red line). The downward triangle indicates the retention time of  $m/z$  216 (BMPO-OH). (b) MS spectrum of  $[\text{BMPO} + \text{H}]^+$  with nominal  $m/z$  200. (c)  $\text{MS}^2$  spectrum of  $m/z$  200, with the characteristic fragment ion  $m/z$  144.0639 ( $[\text{BMPO} + \text{H}]^+ - m/z$  56.0626). (d) Proposed fragmentation pathway for  $m/z$  200. The most abundant fragment ion present in (c) corresponds to the loss of  $\text{C}_4\text{H}_8$  from  $[\text{BMPO} + \text{H}]^+$ . (e) MS spectrum of  $[\text{BMPO-OH}]^+$  with  $m/z$  216. (f) The  $\text{MS}^2$  spectrum of  $m/z$  216, with the characteristic fragment ion  $m/z$  160.0590 ( $[\text{BMPO-OH}]^+ - m/z$  56.0624). (g) Proposed fragmentation pathway for  $m/z$  216. The observed loss of  $\text{C}_4\text{H}_8$  is characteristic of the fragmentation of the *t*-butoxycarbonyl function of BMPO.

tions within 0.1 min, followed by column re-equilibration for 9.9 min before the next run. The ESI-Q-TOF instrument was operated in the positive ionization mode (ESI+) with an ionization voltage of 1900 V. Fragmentation of protonated ions was conducted using the automatic MS/MS mode. Spectra were recorded over the mass range of  $m/z$  100–3000. Data analysis was performed using the qualitative data analysis software (Rev. B. 06.00, Agilent).

## 2.4 Kinetic modelling


The chemical reactions used to describe the BMPO/SOA/ $\text{Fe}^{2+}/\text{H}_2\text{O}$  system, including Fenton-like reactions, are listed along with their rate coefficients in Table S1. From this set of 25 reactions, 16 were optimized using the MCGA method and parameter ranges are given in Table S1 to illustrate the uncertainty arising from global optimization. For all other parameters reference values were taken from the literature, which remained fixed during optimization. Kinetic rate coefficients of a large set of chemical reactions were determined using a uniformly sampled Monte Carlo search seeding a genetic algorithm (MCGA method; Berkemeier et al., 2013; Arangio et al., 2015) as the global optimization method. This algorithm optimizes the

correlation between a kinetic model and experimental data in order to constrain the input parameters of the model. Genetic algorithms mimic processes known from natural evolution and offer mechanisms such as crossover and mutation to avoid convergence towards local minima. In the kinetic model, ROOH represents all organic hydroperoxides without resolving individual structures. This is a simplification, which is necessary for the kinetic modelling but seems to return consistent results.

## 3 Results and discussion

Figure 2 indicates that EPR spectra of laboratory generated SOA by  $\alpha$ -pinene (spectrum a),  $\beta$ -pinene (spectrum b), limonene (spectrum c) and isoprene (spectrum d) SOA were composed of four major peaks, whereas naphthalene SOA (spectrum e) exhibited no significant signals. These four peaks were also found for field samples (spectrum f) and became more prominent in the presence of  $\text{Fe}^{2+}$  (spectrum g). In addition, the same splitting was also observed in a solution of *tert*-butyl hydroperoxide (spectrum h). Four-line signals generated by hyperfine splittings are characteristic of BMPO-trapped OH radicals in water solution,






**Figure 4.** OH formation efficiency by SOA. (a) Concentrations of OH radicals formed in water extracts of SOA of  $\beta$ -pinene (black),  $\alpha$ -pinene (blue), limonene (red), isoprene (purple) and naphthalene (pink) as a function of SOA concentrations in the aqueous phase. The formation efficiency of OH (molar concentration ratio of OH to SOA:  $[\text{BMPO-OH}]/[\text{SOA}]$ , in %) in iron containing SOA water extracts against molar concentration ratios of  $\text{FeSO}_4$  and SOA ( $[\text{Fe}^{2+}]/[\text{SOA}]$ ) by (b)  $\beta$ -pinene, (c)  $\alpha$ -pinene and (d) limonene. The markers are experimental data, and the solid curves with shaded area are modelled with uncertainty.

as shown in the spectrum (spectrum i) for solutions of  $\text{H}_2\text{O}_2$  and  $\text{Fe}^{2+}$ , generating OH via the Fenton reaction ( $\text{Fe}^{2+} + \text{H}_2\text{O}_2 \rightarrow \text{Fe}^{3+} + \text{OH}^- + \cdot\text{OH}$  (Zhao et al., 2001).

Figure 3 shows LC-MS chromatograms of the BMPO-OH adduct ( $m/z$  216.121) for aqueous BMPO solutions (black line) and for BMPO in aqueous  $\beta$ -pinene SOA extract (red line). A strong peak is observed at a retention time of 11.6 min for BMPO in aqueous  $\beta$ -pinene SOA extract, but not for the aqueous BMPO solution, which served as a blank. Confirmation of the BMPO structure for  $m/z$  216.121 was achieved by comparing MS<sup>2</sup> spectra of  $[\text{BMPO} + \text{H}^+]^+$  ( $m/z$  200.126) from the aqueous standard and  $m/z$  216.121. In both cases the loss of a characteristic fragment with a mass of 56.062 Da is observed (panel c and f), which corresponds to the loss of  $\text{C}_4\text{H}_8$  from the *t*-butoxycarbonyl function of BMPO. The above LC-MS/MS analysis confirms the presence of OH radicals in  $\beta$ -pinene SOA extracts observed by EPR shown in Fig. 2.

The EPR and LC-MS/MS observations provide strong evidence that OH radicals are generated in water extracts of SOA by  $\alpha$ -pinene,  $\beta$ -pinene, limonene and isoprene as well as field fine particles, which can be enhanced by  $\text{Fe}^{2+}$ . Note



**Figure 5.** OH yield of  $\beta$ -pinene SOA in three different kinds of pure water: Milli-Q (squares), Savillex (triangles) and TraceSELECT (Sigma, crosses).

that additional hyperfine splitting is observed for monoterpene and isoprene SOA and especially for field samples, indicating the presence of organic radicals. Figure 4a shows that the amount of OH radicals trapped by BMPO increases as the SOA concentration increases in the aqueous phase. The OH yield from  $\beta$ -pinene SOA is the highest generating  $\sim 1.5 \mu\text{M}$  of OH radicals at 1.5 mM SOA concentration, followed by  $\alpha$ -pinene, isoprene and limonene SOA. Naphthalene SOA has a negligible yield of OH radicals.


For assessment of potential interferences from trace amounts of impurities such as transition metals in water, the OH yield was also measured in water with three different purity grades – Milli-Q water (18.2 M, Thermo Scientific<sup>TM</sup> Barnstead<sup>TM</sup> GenPure<sup>TM</sup> xCAD Plus ultrapure water system), TraceSELECT<sup>®</sup> Ultra ACS reagent water (Sigma Aldrich) and Savillex water (DST-1000 Acid Purification System) – which results in excellent agreement (Fig. 5) confirming that OH radicals can be formed in the absence of transition metals.

Ambient particulate matter is often associated with iron ions, which play an important role in aerosol chemistry via Fenton-like reactions (Deguillaume et al., 2005). To investigate the effects of transition metals on OH formation by SOA, different concentrations of  $\text{Fe}^{2+}$  were added in SOA water extracts. Figure 4b–d show the OH formation efficiency (molar concentration ratio of OH and SOA:  $[\text{BMPO-OH}]/[\text{SOA}]$ , in %) of  $\beta$ -pinene,  $\alpha$ -pinene and limonene SOA as a function of molar concentration ratio of  $\text{FeSO}_4$  to SOA ( $[\text{Fe}^{2+}]/[\text{SOA}]$ ). The OH formation efficiency reaches maximum values of 1.5 % for  $\beta$ -pinene SOA, 1.1 % for  $\alpha$ -pinene SOA and 0.5 % for limonene. Different behaviours in OH formation efficiency of limonene compared to  $\alpha$ -pinene and  $\beta$ -pinene may be induced by different organic hydroperoxide concentrations and different R subgroup structure of

ROOH. This order is the same as the order of the relative contribution of organic peroxides in these types of SOA (Docherty et al., 2005). For isoprene SOA, the first results of ongoing experiments indicate a significant increase of OH yield with increasing  $\text{Fe}^{2+}$  concentrations. The EPR spectra of the isoprene SOA show a dependence on the oxidant concentration level in the PAM chamber. The more complex behaviour of the isoprene SOA from OH photooxidation is under investigation and will be presented in a follow-up study.

The observed formation of OH radicals is most likely due to hydrolysis and thermal decomposition of organic hydroperoxides (ROOH), which account for the predominant fraction of terpene SOA (Docherty et al., 2005; Epstein et al., 2014) as well as in rain water (Hellpointner and Gäb, 1989), but they have little contribution for naphthalene SOA (Kautzman et al., 2010). ROOH are formed via multigenerational gas-phase oxidation and autooxidation, introducing multiple hydroperoxy functional groups forming extremely low volatility organic compounds (Crounse et al., 2013; Ehn et al., 2014). Due to the low binding energy of the O–O bond induced by the electron-donating R group, ROOH are well-known to undergo thermal homolytic cleavage ( $\text{ROOH} \rightarrow \text{RO}^\bullet + \bullet\text{OH}$ ; Nam et al., 2000). In the presence of  $\text{Fe}^{2+}$ , it has been reported that decomposition of ROOH can be enhanced mainly via Fenton-like reactions leading to heterolytic cleavage of the O–O bond in the following two ways depending on the pH and reaction environments:  $\text{ROOH} + \text{Fe}^{2+} \rightarrow \text{RO}^\bullet + \text{OH}^- + \text{Fe}^{3+}$  or  $\text{ROOH} + \text{Fe}^{2+} \rightarrow \text{RO}^- + \bullet\text{OH} + \text{Fe}^{3+}$  (Goldstein and Meyerstein, 1999; Deguillaume et al., 2005). Note that homolytic cleavage can be catalysed by iron ions (Foster and Caradonna, 2003). The formed alkoxy radicals ( $\text{RO}^\bullet$ ) were trapped by BMPO and found to increase as the  $\text{Fe}^{2+}$  concentration increases (Fig. 6). The formation of organic radicals in  $\alpha$ -pinene and limonene SOA has been also detected in the previous studies (Pavlovic and Hopke, 2010; Chen et al., 2011). As shown in Fig. 4, the chemical box model including the above three ROOH decomposition pathways reproduces experimental data very well, strongly suggesting that the source of OH radicals is decomposition of ROOH. The decrease of OH radical production with increasing  $\text{Fe}^{2+}$  concentration is supposedly induced by reaction of the BMPO–OH adduct with  $\text{Fe}^{2+}$  (Yamazaki and Piette, 1990) (see also Supplement).

It has been suggested that hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) can be generated from  $\alpha$ - and  $\beta$ -pinene SOA in water, but the mass yield of  $\text{H}_2\text{O}_2$  is  $\sim 0.2\%$  (Wang et al., 2011). In the presence of  $\text{Fe}^{2+}$ ,  $\text{H}_2\text{O}_2$  can yield OH radicals via the Fenton reaction, and the formation efficiency of BMPO–OH adduct by mixtures of  $\text{H}_2\text{O}_2$  with  $\text{Fe}^{2+}$  was measured to be  $\sim 0.6\%$  (Fig. S2). Thus, the potential contribution of generated  $\text{H}_2\text{O}_2$  to OH yields in  $\beta$ - and  $\alpha$ -pinene SOA extracts is much lower than the observed OH radicals. Moreover, the OH yield was not affected, even if  $\beta$ -pinene SOA was dried under a  $\text{N}_2$  flow before the water extraction to evaporate particle-phase




**Figure 6.** Formation efficiency of organic radicals. Molar concentration ratio of organic radicals to SOA ( $[\text{BMPO-OR}]/[\text{SOA}]$ , in %) in mixtures of  $\text{Fe}^{2+}$  and SOA solutions.

$\text{H}_2\text{O}_2$ . Hence it is clear that the  $\text{H}_2\text{O}_2$  in SOA should not be the dominant source of OH radicals observed in this study.


#### 4 Implications

The implications of this finding are illustrated in Figs. 7 and 8. The orange area in Fig. 7a shows OH production rate by Fenton reactions between  $\text{Fe}^{2+}$  and  $\text{H}_2\text{O}_2$  forming OH radicals as a function of  $\text{H}_2\text{O}_2$  concentration with typical dissolved iron concentrations in cloud droplets of  $0.1\text{--}2.5\ \mu\text{M}$  (Deguillaume et al., 2005). The green area shows the OH production rate by SOA decomposition in cloud or fog droplets, which ranges of  $\sim 0.01\text{--}100\ \text{nM s}^{-1}$  depending on SOA precursors and the  $\text{Fe}^{2+}$  and SOA concentrations (see Supplement). It clearly shows that SOA decomposition is comparably important to the Fenton reaction in most conditions and that SOA can be the main source of OH radicals at low concentrations of  $\text{H}_2\text{O}_2$  and  $\text{Fe}^{2+}$ . Water-soluble gases such as aldehydes taken up by deliquesced particles may undergo reactions in the presence of OH radicals to form low-volatility products, including organic acids, peroxides, peroxyhemiacetals and oligomers (Lim et al., 2010; Ervens et al., 2011; Liu et al., 2012; Ervens, 2015; Lim and Turpin, 2015; McNeill, 2015). Thus, the formed OH radicals would promote chemical aging of SOA especially in the presence of iron ions (e.g. SOA-coated mineral dust particles) (Chu et al., 2014) and may also induce aqueous-phase oxidation of sulfur dioxide forming sulfuric acid (Harris et al., 2013).

Recent studies have shown that OH radicals can trigger autoxidation reactions in the gas phase, generating highly oxidized and extremely low volatility compounds (Crounse et al., 2013; Ehn et al., 2014). In addition, it has been shown that some radicals can be long-lived in the condensed phase



**Figure 7.** OH production rate in cloud droplets and lung lining fluid. (a) The OH production rate in cloud droplets by SOA decomposition compared to the classical Fenton reaction. The data points were measured in the absence of  $\text{Fe}^{2+}$  for different precursors of  $\beta$ -pinene (black squares),  $\alpha$ -pinene (blue circles), limonene (red upward triangles) and isoprene (purple downward triangles). The shaded green area represents the possible range in the presence of iron as a function of SOA concentration in the aqueous phase, which is based on the minimum and maximum OH radical production efficiency of SOA in Fig. 4. The dashed lines represent OH production rates due to the Fenton reaction from  $\text{H}_2\text{O}_2$  with typical dissolved iron concentrations ( $\text{Fe}^{2+} : \text{Fe}^{3+} = 1 : 1$ ) of 0.1 and 2.5  $\mu\text{M}$ . (b) The OH production rate in lung lining fluid by SOA decomposition as a function of ambient SOA concentrations, and by the classical Fenton reaction as a function of  $\text{H}_2\text{O}_2$  concentrations with typical dissolved iron concentrations ( $\text{Fe}^{2+} : \text{Fe}^{3+} = 1 : 1$ ) of 100 and 1 nM. The purple shaded area represents patients with respiratory disease exhibiting high  $\text{H}_2\text{O}_2$  concentrations in the bronchoalveolar lavage (Corradi et al., 2008).



**Figure 8.** Implications of OH formation by SOA. Formation of OH radicals upon decomposition of organic hydroperoxides (ROOH) in secondary organic aerosol leads to rapid chemical aging of SOA particles upon deliquescence and cloud or fog processing in the atmosphere as well as oxidative stress upon inhalation and deposition in the human respiratory tract. Mixing and Fenton-like reactions of iron with ROOH from SOA can occur both in atmospheric particles and in the lung lining fluid.

(Shiraiwa et al., 2011b; Gehling and Dellinger, 2013) by interacting with transition metals (Truong et al., 2010). We hypothesize that OH radicals formed from SOA decomposition could also trigger autoxidation in the condensed phase. Such

a self-amplification cycle of SOA formation and aging may be relevant for example in the Amazon, where cloud and fog processing are important pathways forming a high fraction of SOA with high O : C ratio, resulting in an enhancement of cloud condensation nuclei activity of particles (Pöschl et al., 2010; Pöhlker et al., 2012). Organic peroxides are often used as the agent of the vulcanization processes to initiate the radical polymerization by forming free radicals, which abstract hydrogen atoms from the elastomer molecules converting them into radicals that undergo oligomerization to form elastic polymer or rubber. Similar processes might also occur in SOA particles (“SOA vulcanization”), which may contribute to formation of dimers and oligomers observed in SOA particles (Kalberer et al., 2004) possibly leading to the occurrence of an amorphous solid state (Virtanen et al., 2010; Koop et al., 2011; Shiraiwa et al., 2011a; Renbaum-Wolff et al., 2013; Kidd et al., 2014).

In indoor air, terpenes are commonly found at higher concentrations than in the ambient air due to their widespread use as solvents and odorants in cleaning products and air fresheners (Weschler, 2011). Depending on precursor concentrations, the SOA concentration in indoor air can reach up to  $30 \mu\text{g m}^{-3}$  with the highest contribution from limonene SOA (Waring, 2014). To evaluate potential adverse health effects by SOA deposition into the lungs, we estimated the OH production rate by SOA within the lung lining fluid (LLF) as

a function of ambient SOA concentration considering breathing and deposition rates (see Supplement) (Fig. 7b). The pH of lung lining fluid for healthy people is about 7.4. Our recent experiments have shown that the formation of OH radicals was increased by  $\sim 20\%$  at a pH of 7.4 in a phosphate-buffered saline solution. Thus, the OH production rate by SOA decomposition shown in Fig. 7b may represent the lower limit. We intend to investigate pH effects on OH formation in detail in follow-up studies.

Figure 7b also shows the OH production rate by the Fenton reaction with typical iron (Gutteridge et al., 1996) and  $\text{H}_2\text{O}_2$  concentrations in the LLF (Corradi et al., 2008). Patients with respiratory diseases are reported to have high  $\text{H}_2\text{O}_2$  concentrations in the bronchoalveolar lavage (Corradi et al., 2008) (as shown in shaded purple area), and the Fenton reaction may be the main source of OH radicals for such patients. However, for healthy people with low  $\text{H}_2\text{O}_2$  and  $\text{Fe}^{2+}$  concentrations, SOA decomposition can be more important than the Fenton process under high ambient or indoor SOA concentrations. Excess concentrations of reactive oxygen species including hydrogen peroxide, OH radicals (and potentially also organic radicals) are shown to cause oxidative stress to human lung fibroblasts, alveolar cells and tissues (Pöschl and Shiraiwa, 2015). Thus, in polluted indoor or urban megacities with high SOA concentration such as in Beijing, SOA particles may play a critical role in adverse aerosol health effects.

**The Supplement related to this article is available online at doi:10.5194/acp-16-1761-2016-supplement.**

**Acknowledgements.** This work was funded by the Max Planck Society. C. J. Kampf acknowledges financial support by the German Research Foundation (DFG project KA 4008/1-1).

The article processing charges for this open-access publication were covered by the Max Planck Society.

Edited by: F. Keutsch

## References

- Anglada, J. M., Martins-Costa, M., Ruiz-López, M. F., and Francisco, J. S.: Spectroscopic signatures of ozone at the air–water interface and photochemistry implications, *P. Natl. Acad. Sci. USA*, 111, 11618–11623, 2014.
- Arakaki, T., Anastasio, C., Kuroki, Y., Nakajima, H., Okada, K., Kotani, Y., Handa, D., Azechi, S., Kimura, T., and Tsuchioka, A.: A general scavenging rate constant for reaction of hydroxyl radical with organic carbon in atmospheric waters, *Environ. Sci. Technol.*, 47, 8196–8203, 2013.
- Arangio, A. M., Slade, J. H., Berkemeier, T., Pöschl, U., Knopf, D. A., and Shiraiwa, M.: Multiphase Chemical Kinetics of OH Radical Uptake by Molecular Organic Markers of Biomass Burning Aerosols: Humidity and Temperature Dependence, Surface Reaction, and Bulk Diffusion, *J. Phys. Chem. A*, 119, 4533–4544, 2015.
- Badali, K. M., Zhou, S., Aljawhary, D., Antiñolo, M., Chen, W. J., Lok, A., Mungall, E., Wong, J. P. S., Zhao, R., and Abbatt, J. P. D.: Formation of hydroxyl radicals from photolysis of secondary organic aerosol material, *Atmos. Chem. Phys.*, 15, 7831–7840, doi:10.5194/acp-15-7831-2015, 2015.
- Berkemeier, T., Huisman, A. J., Ammann, M., Shiraiwa, M., Koop, T., and Pöschl, U.: Kinetic regimes and limiting cases of gas uptake and heterogeneous reactions in atmospheric aerosols and clouds: a general classification scheme, *Atmos. Chem. Phys.*, 13, 6663–6686, doi:10.5194/acp-13-6663-2013, 2013.
- Bruns, E. A., El Haddad, I., Keller, A., Klein, F., Kumar, N. K., Pieber, S. M., Corbin, J. C., Slowik, J. G., Brune, W. H., Baltensperger, U., and Prévôt, A. S. H.: Inter-comparison of laboratory smog chamber and flow reactor systems on organic aerosol yield and composition, *Atmos. Meas. Tech.*, 8, 2315–2332, doi:10.5194/amt-8-2315-2015, 2015.
- Chen, X. and Hopke, P.: A chamber study of secondary organic aerosol formation by limonene ozonolysis, *Indoor air*, 20, 320–328, 2010.
- Chen, X., Hopke, P. K., and Carter, W. P.: Secondary organic aerosol from ozonolysis of biogenic volatile organic compounds: chamber studies of particle and reactive oxygen species formation, *Environ. Sci. Technol.*, 45, 276–282, 2010.
- Chen, X., Hopke, P. K., and Carter, W. P. L.: Secondary Organic Aerosol from Ozonolysis of Biogenic Volatile Organic Compounds: Chamber Studies of Particle and Reactive Oxygen Species Formation, *Environ. Sci. Technol.*, 45, 276–282, 2011.
- Chu, B., Liu, Y., Li, J., Takekawa, H., Liggio, J., Li, S.-M., Jiang, J., Hao, J., and He, H.: Decreasing effect and mechanism of  $\text{FeSO}_4$  seed particles on secondary organic aerosol in  $\alpha$ -pinene photooxidation, *Environ. Pollut.*, 193, 88–93, 2014.
- Corradi, M., Pignatti, P., Brunetti, G., Goldoni, M., Caglieri, A., Nava, S., Moscato, G., and Balbi, B.: Comparison between exhaled and bronchoalveolar lavage levels of hydrogen peroxide in patients with diffuse interstitial lung diseases, *Acta Biomed.*, 79, 73–78, 2008.
- Crounse, J. D., Nielsen, L. B., Jørgensen, S., Kjaergaard, H. G., and Wennberg, P. O.: Autoxidation of organic compounds in the atmosphere, *J. Phys. Chem. Lett.*, 4, 3513–3520, 2013.
- Deguillaume, L., Leriche, M., Desboeufs, K., Mailhot, G., George, C., and Chaumerliac, N.: Transition Metals in Atmospheric Liquid Phases: Sources, Reactivity, and Sensitive Parameters, *Chem. Rev.*, 105, 3388–3431, 2005.
- Docherty, K. S., Wu, W., Lim, Y. B., and Ziemann, P. J.: Contributions of organic peroxides to secondary aerosol formed from reactions of monoterpenes with  $\text{O}_3$ , *Environ. Sci. Technol.*, 39, 4049–4059, 2005.
- Donahue, N. M., Henry, K. M., Mentel, T. F., Kiendler-Scharr, A., Spindler, C., Bohn, B., Brauers, T., Dorn, H. P., Fuchs, H., Tillmann, R., Wahner, A., Saathoff, H., Naumann, K.-H., Möhler, O., Leisner, T., Müller, L., Reinnig, M.-C., Hoffmann, T., Salo, K., Hallquist, M., Frosch, M., Bilde, M., Tritscher, T., Barmet, P., Praplan, A. P., DeCarlo, P. F., Dommen, J., Prévôt, A. S. H., and



- Baltensperger, U.: Aging of biogenic secondary organic aerosol via gas-phase OH radical reactions, *P. Natl. Acad. Sci. USA*, 109, 13503–13508, 2012.
- Ehn, M., Thornton, J. A., Kleist, E., Sipila, M., Junninen, H., Pullinen, I., Springer, M., Rubach, F., Tillmann, R., Lee, B., Lopez-Hilfiker, F., Andres, S., Acir, I.-H., Rissanen, M., Jokinen, T., Schobesberger, S., Kangasluoma, J., Kontkanen, J., Nieminen, T., Kurten, T., Nielsen, L. B., Jorgensen, S., Kjaergaard, H. G., Canagaratna, M., Dal Maso, M., Berndt, T., Petaja, T., Wahner, A., Kerminen, V.-M., Kulmala, M., Worsnop, D. R., Wildt, J., and Mentel, T. F.: A large source of low-volatility secondary organic aerosol, *Nature*, 506, 476–479, 2014.
- Epstein, S. A., Blair, S. L., and Nizkorodov, S. A.: Direct photolysis of  $\alpha$ -pinene ozonolysis secondary organic aerosol: effect on particle mass and peroxide content, *Environ. Sci. Technol.*, 48, 11251–11258, 2014.
- Ervens, B.: Modeling the Processing of Aerosol and Trace Gases in Clouds and Fogs, *Chem. Rev.*, 115, 4157–4198, 2015.
- Ervens, B., Turpin, B. J., and Weber, R. J.: Secondary organic aerosol formation in cloud droplets and aqueous particles (aqSOA): a review of laboratory, field and model studies, *Atmos. Chem. Phys.*, 11, 11069–11102, doi:10.5194/acp-11-11069-2011, 2011.
- Faber, P., Drewnick, F., Veres, P. R., Williams, J., and Borrmann, S.: Anthropogenic sources of aerosol particles in a football stadium: Real-time characterization of emissions from cigarette smoking, cooking, hand flares, and color smoke bombs by high-resolution aerosol mass spectrometry, *Atmos. Environ.*, 77, 1043–1051, 2013.
- Foster, T. L. and Caradonna, J. P.:  $\text{Fe}^{2+}$ -catalyzed heterolytic RO-OH bond cleavage and substrate oxidation: A functional synthetic non-heme iron monooxygenase system, *J. Am. Chem. Soc.*, 125, 3678–3679, 2003.
- Fuller, S., Wragg, F., Nutter, J., and Kalberer, M.: Comparison of on-line and off-line methods to quantify reactive oxygen species (ROS) in atmospheric aerosols, *Atmos. Environ.*, 92, 97–103, 2014.
- Gehling, W. and Dellinger, B.: Environmentally Persistent Free Radicals and Their Lifetimes in  $\text{PM}_{2.5}$ , *Environ. Sci. Technol.*, 47, 8172–8178, 2013.
- Goldstein, S. and Meyerstein, D.: Comments on the mechanism of the “Fenton-like” reaction, *Acc. Chem. Res.*, 32, 547–550, 1999.
- Gutteridge, J. M. C., Mumby, S., Quinlan, G. J., Chung, K. F., and Evans, T. W.: Pro-oxidant iron is present in human pulmonary epithelial lining fluid: implications for oxidative stress in the lung, *Biochem. Biophys. Res. Commun.*, 220, 1024–1027, 1996.
- Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, Th. F., Monod, A., Prévôt, A. S. H., Seinfeld, J. H., Surratt, J. D., Szmigielski, R., and Wildt, J.: The formation, properties and impact of secondary organic aerosol: current and emerging issues, *Atmos. Chem. Phys.*, 9, 5155–5236, doi:10.5194/acp-9-5155-2009, 2009.
- Harris, E., Sinha, B., van Pinxteren, D., Tilgner, A., Fomba, K. W., Schneider, J., Roth, A., Gnauk, T., Fahlbusch, B., Mertes, S., Lee, T., Collett, J., Foley, S., Borrmann, S., Hoppe, P., and Herrmann, H.: Enhanced Role of Transition Metal Ion Catalysis During In-Cloud Oxidation of  $\text{SO}_2$ , *Science*, 340, 727–730, 2013.
- Hellpointner, E. and Gäh, S.: Detection of methyl, hydroxymethyl and hydroxyethyl hydroperoxides in air and precipitation, *Nature*, 337, 631–634, 1989.
- Herrmann, H., Tilgner, A., Barzagli, P., Majdik, Z., Gligorovski, S., Poulain, L., and Monod, A.: Towards a more detailed description of tropospheric aqueous phase organic chemistry: CAPRAM 3.0, *Atmos. Environ.*, 39, 4351–4363, 2005.
- Huang, R.-J., Zhang, Y., Bozzetti, C., Ho, K.-F., Cao, J.-J., Han, Y., Daellenbach, K. R., Slowik, J. G., Platt, S. M., Canonaco, F., Zotter, P., Wolf, R., Pieber, S. M., Bruns, E. A., Crippa, M., Ciarelli, G., Piazzalunga, A., Schwikowski, M., Abbaszade, G., Schnelle-Kreis, J., Zimmermann, R., An, Z., Szidat, S., Baltensperger, U., Haddad, I. E., and Prevot, A. S. H.: High secondary aerosol contribution to particulate pollution during haze events in China, *Nature*, 514, 218–222, 2014.
- Jacob, D. J.: Chemistry of OH in remote clouds and its role in the production of formic acid and peroxyxymonosulfate, *J. Geophys. Res.-Atmos.*, 91, 9807–9826, 1986.
- Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H., DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M., Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., Dunlea, E. J., Huffman, J. A., Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimono, A., Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel, J. R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L. R., Wood, E. C., Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of organic aerosols in the atmosphere, *Science*, 326, 1525–1529, 2009.
- Jokinen, T., Berndt, T., Makkonen, R., Kerminen, V.-M., Junninen, H., Paasonen, P., Stratmann, F., Herrmann, H., Guenther, A. B., Worsnop, D. R., Kulmala, M., Ehn, M., and Sipilä, M.: Production of extremely low volatile organic compounds from biogenic emissions: Measured yields and atmospheric implications, *P. Natl. Acad. Sci. USA*, 112, 7123–7128, 2015.
- Kalberer, M., Paulsen, D., Sax, M., Steinbacher, M., Dommen, J., Prevot, A., Fisseha, R., Weingartner, E., Frankevich, V., and Zenobi, R.: Identification of polymers as major components of atmospheric organic aerosols, *Science*, 303, 1659–1662, 2004.
- Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., Van Dingenen, R., Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P., Balkanski, Y., Fuzzi, S., Horth, J., Moortgat, G. K., Winterhalter, R., Myhre, C. E. L., Tsigaridis, K., Vignati, E., Stephanou, E. G., and Wilson, J.: Organic aerosol and global climate modelling: a review, *Atmos. Chem. Phys.*, 5, 1053–1123, doi:10.5194/acp-5-1053-2005, 2005.
- Kang, E., Root, M. J., Toohey, D. W., and Brune, W. H.: Introducing the concept of Potential Aerosol Mass (PAM), *Atmos. Chem. Phys.*, 7, 5727–5744, doi:10.5194/acp-7-5727-2007, 2007.
- Kautzman, K. E., Surratt, J. D., Chan, M. N., Chan, A. W. H., Hersey, S. P., Chhabra, P. S., Dalleska, N. F., Wennberg, P. O.,

- Flagan, R. C., and Seinfeld, J. H.: Chemical Composition of Gas- and Aerosol-Phase Products from the Photooxidation of Naphthalene, *J. Phys. Chem. A*, 114, 913–934, 2010.
- Kidd, C., Perraud, V., Wingen, L. M., and Finlayson-Pitts, B. J.: Integrating phase and composition of secondary organic aerosol from the ozonolysis of alpha-pinene, *P. Natl. Acad. Sci. USA*, 111, 7552–7557, 2014.
- Koop, T., Bookhold, J., Shiraiwa, M., and Pöschl, U.: Glass transition and phase state of organic compounds: dependency on molecular properties and implications for secondary organic aerosols in the atmosphere, *Phys. Chem. Chem. Phys.*, 13, 19238–19255, 2011.
- Lambe, A. T., Ahern, A. T., Williams, L. R., Slowik, J. G., Wong, J. P. S., Abbatt, J. P. D., Brune, W. H., Ng, N. L., Wright, J. P., Croasdale, D. R., Worsnop, D. R., Davidovits, P., and Onasch, T. B.: Characterization of aerosol photooxidation flow reactors: heterogeneous oxidation, secondary organic aerosol formation and cloud condensation nuclei activity measurements, *Atmos. Meas. Tech.*, 4, 445–461, doi:10.5194/amt-4-445-2011, 2011.
- Lambe, A. T., Chhabra, P. S., Onasch, T. B., Brune, W. H., Hunter, J. F., Kroll, J. H., Cummings, M. J., Brogan, J. F., Parmar, Y., Worsnop, D. R., Kolb, C. E., and Davidovits, P.: Effect of oxidant concentration, exposure time, and seed particles on secondary organic aerosol chemical composition and yield, *Atmos. Chem. Phys.*, 15, 3063–3075, doi:10.5194/acp-15-3063-2015, 2015.
- Lim, Y. B. and Turpin, B. J.: Laboratory evidence of organic peroxide and peroxyhemiacetal formation in the aqueous phase and implications for aqueous OH, *Atmos. Chem. Phys.*, 15, 12867–12877, doi:10.5194/acp-15-12867-2015, 2015.
- Lim, Y. B., Tan, Y., Perri, M. J., Seitzinger, S. P., and Turpin, B. J.: Aqueous chemistry and its role in secondary organic aerosol (SOA) formation, *Atmos. Chem. Phys.*, 10, 10521–10539, doi:10.5194/acp-10-10521-2010, 2010.
- Liu, Y., Monod, A., Tritscher, T., Praplan, A. P., DeCarlo, P. F., Temime-Roussel, B., Quivet, E., Marchand, N., Dommen, J., and Baltensperger, U.: Aqueous phase processing of secondary organic aerosol from isoprene photooxidation, *Atmos. Chem. Phys.*, 12, 5879–5895, doi:10.5194/acp-12-5879-2012, 2012.
- McNeill, V. F.: Aqueous organic chemistry in the atmosphere: Sources and chemical processing of organic aerosols, *Environ. Sci. Technol.*, 49, 1237–1244, 2015.
- Mentel, T. F., Springer, M., Ehn, M., Kleist, E., Pullinen, I., Kurtén, T., Rissanen, M., Wahner, A., and Wildt, J.: Formation of highly oxidized multifunctional compounds: autoxidation of peroxy radicals formed in the ozonolysis of alkenes – deduced from structure–product relationships, *Atmos. Chem. Phys.*, 15, 6745–6765, doi:10.5194/acp-15-6745-2015, 2015.
- Nam, W., Han, H. J., Oh, S.-Y., Lee, Y. J., Choi, M.-H., Han, S.-Y., Kim, C., Woo, S. K., and Shin, W.: New insights into the mechanisms of OO bond cleavage of hydrogen peroxide and tert-alkyl hydroperoxides by iron (III) porphyrin complexes, *J. Am. Chem. Soc.*, 122, 8677–8684, 2000.
- Ortega, A. M., Hayes, P. L., Peng, Z., Palm, B. B., Hu, W., Day, D. A., Li, R., Cubison, M. J., Brune, W. H., Graus, M., Warneke, C., Gilman, J. B., Kuster, W. C., de Gouw, J. A., and Jimenez, J. L.: Real-time measurements of secondary organic aerosol formation and aging from ambient air in an oxidation flow reactor in the Los Angeles area, *Atmos. Chem. Phys. Discuss.*, 15, 21907–21958, doi:10.5194/acpd-15-21907-2015, 2015.
- Pavlovic, J. and Hopke, P. K.: Detection of radical species formed by the ozonolysis of  $\alpha$ -pinene, *J. Atmos. Chem.*, 66, 137–155, 2010.
- Pöhlker, C., Wiedemann, K. T., Sinha, B., Shiraiwa, M., Gunthe, S. S., Smith, M., Su, H., Artaxo, P., Chen, Q., Cheng, Y., Elbert, W., Gilles, M. K., Kilcoyne, A. L. D., Moffet, R. C., Weigand, M., Martin, S. T., Pöschl, U., and Andreae, M. O.: Biogenic potassium salt particles as seeds for secondary organic aerosol in the Amazon, *Science*, 337, 1075–1078, 2012.
- Pöschl, U. and Shiraiwa, M.: Multiphase Chemistry at the Atmosphere–Biosphere Interface Influencing Climate and Public Health in the Anthropocene, *Chem. Rev.*, 115, 4440–4475, 2015.
- Pöschl, U., Martin, S. T., Sinha, B., Chen, Q., Gunthe, S. S., Huffman, J. A., Borrmann, S., Farmer, D. K., Garland, R. M., Helas, G., Jimenez, J. L., King, S. M., Manzi, A., Mikhailov, E., Pauliquevis, T., Petters, M. D., Prenni, A. J., Roldin, P., Rose, D., Schneider, J., Su, H., Zorn, S. R., Artaxo, P., and Andreae, M. O.: Rainforest Aerosols as Biogenic Nuclei of Clouds and Precipitation in the Amazon, *Science*, 329, 1513–1516, 2010.
- Renbaum-Wolff, L., Grayson, J. W., Bateman, A. P., Kuwata, K., Sellier, M., Murray, B. J., Schilling, J. E., Martin, S. T., and Bertram, A. K.: Viscosity of  $\alpha$ -pinene secondary organic material and implications for particle growth and reactivity, *P. Natl. Acad. Sci. USA*, 110, 8014–8019, 2013.
- Shiraiwa, M., Ammann, M., Koop, T., and Pöschl, U.: Gas uptake and chemical aging of semisolid organic aerosol particles, *P. Natl. Acad. Sci. USA*, 108, 11003–11008, 2011a.
- Shiraiwa, M., Sosedova, Y., Rouvière, A., Yang, H., Zhang, Y., Abbatt, J. P., Ammann, M., and Pöschl, U.: The role of long-lived reactive oxygen intermediates in the reaction of ozone with aerosol particles, *Nat. Chem.*, 3, 291–295, 2011b.
- Shiraiwa, M., Yee, L. D., Schilling, K. A., Loza, C. L., Craven, J. S., Zuend, A., Ziemann, P. J., and Seinfeld, J. H.: Size distribution dynamics reveal particle-phase chemistry in organic aerosol formation, *P. Natl. Acad. Sci. USA*, 110, 11746–11750, 2013.
- Solomon, S.: Climate change 2007-the physical science basis: Working group I contribution to the fourth assessment report of the IPCC, Cambridge University Press, Cambridge, 2007.
- Stoll, S. and Schweiger, A.: EasySpin, a comprehensive software package for spectral simulation and analysis in EPR, *J. Magn. Reson.*, 178, 42–55, 2006.
- Truong, H., Lomnicki, S., and Dellinger, B.: Potential for misidentification of environmentally persistent free radicals as molecular pollutants in particulate matter, *Environ. Sci. Technol.*, 44, 1933–1939, 2010.
- Venkatachari, P. and Hopke, P. K.: Development and evaluation of a particle-bound reactive oxygen species generator, *J. Aerosol. Sci.*, 39, 168–174, 2008.
- Virtanen, A., Joutsensaari, J., Koop, T., Kannosto, J., YliPirilä, P., Leskinen, J., Mäkelä, J. M., Holopainen, J. K., Pöschl, U., Kulmala, M., Worsnop, D. R., and Laaksonen, A.: An amorphous solid state of biogenic secondary organic aerosol particles, *Nature*, 467, 824–827, 2010.
- Wang, Y., Kim, H., and Paulson, S. E.: Hydrogen peroxide generation from alpha- and beta-pinene and toluene secondary organic aerosols, *Atmos. Environ.*, 45, 3149–3156, 2011.

- Waring, M. S.: Secondary organic aerosol in residences: predicting its fraction of fine particle mass and determinants of formation strength, *Indoor Air*, 24, 376–389, 2014.
- Weber, R. T.: Xenon Data Processing Reference, Bruker BioSpin Corporation: Billerica, MA USA, chapter 14, 1–24, 2012.
- Weschler, C. J.: Chemistry in indoor environments: 20 years of research, *Indoor Air*, 21, 205–218, 2011.
- Yamazaki, I. and Piette, L. H.: ESR spin-trapping studies on the reaction of  $\text{Fe}^{2+}$  ions with  $\text{H}_2\text{O}_2$ -reactive species in oxygen toxicity in biology, *J. Biol. Chem.*, 265, 13589–13594, 1990.
- Zhao, H., Joseph, J., Zhang, H., Karoui, H., and Kalyanaraman, B.: Synthesis and biochemical applications of a solid cyclic nitron spin trap: a relatively superior trap for detecting superoxide anions and glutathionyl radicals, *Free Radic. Biol. Med.*, 31, 599–606, 2001.
- Zhu, B.-Z., Shan, G.-Q., Huang, C.-H., Kalyanaraman, B., Mao, L., and Du, Y.-G.: Metal-independent decomposition of hydroperoxides by halogenated quinones: Detection and identification of a quinone ketoxy radical, *P. Natl. Acad. Sci. USA*, 106, 11466–11471, 2009.