



Conditions for super-adiabatic droplet growth after entrainment mixing

Fan Yang¹, Raymond Shaw¹, and Huiwen Xue²

¹Atmospheric Sciences Program and Department of Physics, Michigan Technological University, Houghton, Michigan, USA

²Department of Atmospheric and Oceanic Sciences, School of Physics, Peking University, Beijing, China

Correspondence to: Raymond Shaw (rashaw@mtu.edu)

Received: 29 January 2016 – Published in Atmos. Chem. Phys. Discuss.: 5 February 2016

Revised: 30 May 2016 – Accepted: 8 July 2016 – Published: 29 July 2016

Abstract. Cloud droplet response to entrainment and mixing between a cloud and its environment is considered, accounting for subsequent droplet growth during adiabatic ascent following a mixing event. The vertical profile for liquid water mixing ratio after a mixing event is derived analytically, allowing the reduction to be predicted from the mixing fraction and from the temperature and humidity for both the cloud and environment. It is derived for the limit of homogeneous mixing. The expression leads to a critical height above the mixing level: at the critical height the cloud droplet radius is the same for both mixed and unmixed parcels, and the critical height is independent of the updraft velocity and mixing fraction. Cloud droplets in a mixed parcel are larger than in an unmixed parcel above the critical height, which we refer to as the “super-adiabatic” growth region. Analytical results are confirmed with a bin microphysics cloud model. Using the model, we explore the effects of updraft velocity, aerosol source in the environmental air, and polydisperse cloud droplets. Results show that the mixed parcel is more likely to reach the super-adiabatic growth region when the environmental air is humid and clean. It is also confirmed that the analytical predictions are matched by the volume-mean cloud droplet radius for polydisperse size distributions. The findings have implications for the origin of large cloud droplets that may contribute to onset of collision-coalescence in warm clouds.

1 Introduction

Warm clouds play an important role for the water cycle and energy balance in the atmosphere. However their formation, development and precipitation processes are still not fully understood (e.g., Beard and Ochs III, 1993). Observations show that warm clouds can precipitate within 20 min (e.g., Laird et al., 2000; Göke et al., 2007). One open question is how small cloud droplets, which are on the order of 10 μm , change to rain drops, usually on order of 1 mm, within such a short time. Because condensation growth is slow for droplet size larger than approximately 20 μm , collision growth is believed to be the most important mechanism for warm cloud precipitation (Pruppacher et al., 1998).

However, collision efficiency is very low for droplets smaller than $r \approx 30 \mu\text{m}$ due to hydrodynamic interaction. For example, Hocking (1959) considered two moving droplets in the Stokes flow approximation and found that the collision efficiency for a $r = 19 \mu\text{m}$ droplet with smaller droplets is mostly less than 0.1, and even for a $r = 30 \mu\text{m}$ droplet it is mostly less than 0.5. Such low collision efficiency suppresses the time required for drizzle and precipitation formation. Therefore, large cloud droplets are needed to efficiently initiate precipitation. There are several hypotheses to explain the formation of large cloud droplets: for example, the stochastic collision process itself may produce a small number of “lucky” droplets with larger growth rates (Kostinski and Shaw, 2005). Another possible mechanism is due to the giant cloud condensation nuclei (CCN). Observational results show that giant and ultragiant CCN often exist in the atmosphere, and simulation results indicate that they can be

sufficient to start the rain precipitation (e.g., Johnson, 1982; Feingold et al., 1999; Yin et al., 2000; Blyth et al., 2003; Jensen and Lee, 2008; Cheng et al., 2009). But in this paper we focus on mechanisms involving the condensation process. For example, results from Lagrangian tracking studies suggest that large droplets from condensation growth within parcels having favored trajectories can trigger collisions and drizzle formation in warm clouds (Lasher-Trapp et al., 2005; Cooper et al., 2013; Magaritz-Ronen et al., 2014, 2016; Naumann and Seifert, 2015; de Lozar and Muesle, 2016). Korolev et al. (2013) proposed that the droplet size distribution can be broadened through diffusion growth due to cloud base mixing and vertical fluctuation. Perhaps counter-intuitively, the mixing and entrainment that occurs during cloud evolution itself may be responsible for generating large cloud droplets (Baker et al., 1980). The possibility that entrainment and subsequent growth can lead to droplets larger than would occur in an unmixed parcel has occupied the attention of the cloud physics community for several decades (e.g., Baker et al., 1980; Jensen et al., 1985; Paluch and Knight, 1986; Su et al., 1998; Cooper et al., 2013; Schmeissner et al., 2015).

Observational results show that the number concentration of cloud droplets at the cloud edge/top is usually smaller than that in the cloud due to entrainment and mixing with environmental air. However, the mean size of cloud droplets at the edge/top might be smaller, equal to, or even larger than that in the cloud (e.g., Burnet and Brenguier, 2007; Lehmann et al., 2009; Lu et al., 2013; Beals et al., 2015), which is thought to be the result of different mixing processes. Baker et al. (1980) proposed two limiting mixing processes: homogeneous and extreme inhomogeneous mixing. Theoretically, mean cloud droplet size will decrease for homogeneous mixing but remains the same for extreme inhomogeneous mixing. However the actual mixing process near the cloud edge/top and the response of cloud droplets to the mixing process are still unclear. Recently, considerable theoretical and computational work has been directed toward understanding the evolution of the droplet size distribution during both homogeneous and inhomogeneous mixing processes (Andrejczuk et al., 2009; Kumar et al., 2014; Tölle and Krueger, 2014; Korolev et al., 2016; Pinsky et al., 2016b, a). Most of these analyses, however, did not consider the subsequent vertical movement of the mixed parcel, which is also relevant to the evolution of cloud droplets (Wang et al., 2009; Yum et al., 2015; Chen et al., 2015). Finally, most theoretical work thus far does not account for the possibility of secondary activation of aerosols after dilution and mixing, although there is compelling experimental evidence that this occurs (Burnet and Brenguier, 2007; Schmeissner et al., 2015).

In this study, we are interested in the change of cloud microphysical properties after isobaric mixing of cloudy and clear-air volumes, assuming the mixing parcel rises adiabatically afterwards. In reality, a cloud parcel can continuously mix with both cloudy air and the environment air through-

out its trajectory. Previous studies (e.g., Cooper et al., 2013; Magaritz-Ronen et al., 2014) have demonstrated some effects of internal mixing, especially due to sedimentation when drizzle is present, and that dilution events often take place repeatedly during parcel ascent. The results presented here do not consider the fully realistic conditions, but instead are purposefully designed so as to avoid the complexity of a real cloud and look at the idealized response to a single dilution event. Our motivating philosophy is that if we can understand the “impulse response” from one mixing event with analytical results, then that understanding can be extended to multiple dilution events. This view of a single mixing event followed by isolated growth is therefore an idealization that allows us to understand the microphysical response in the simplest conditions. We pose the following question: is it possible to achieve “super-adiabatic” droplet diameters as a result of mixing? By super-adiabatic, we mean that the droplet diameter is larger than that observed for an unmixed, closed parcel that grows according to moist-adiabatic conditions (as defined, for example, by Cotton et al. (2011, Chap. 4)). Specifically, we look for the conditions, such as mixing fraction, ambient humidity, aerosol entrainment, secondary activation, and vertical displacement above the mixing level, that influence the ability to produce larger droplets than exist in an unmixed parcel. We first address the problem by deriving analytical results in Sect. 2, and then evaluate the theory and explore conditions for super-adiabatic droplet growth using a microphysical cloud parcel model in Sect. 3. Implications are discussed and results are summarized in Sect. 4.

2 Analytical results

As in previous studies, we consider the final state of the microphysical variables (e.g., liquid water mixing ratio, droplet sizes) after homogeneous mixing (e.g., Korolev et al., 2016). This corresponds to the limit of instantaneous mixing, under which conservation of energy and mass result in a unique dependence of droplet size on the mixing fraction (e.g., Andrejczuk et al., 2006; Burnet and Brenguier, 2007; Gerber et al., 2008; Kumar et al., 2014). Here, we consider the similar two stages of homogeneous mixing process as discussed in Pinsky et al. (2016b), except that the cloud parcel has continuous vertical movement after the mixing event. The first stage is (instantaneous) isobaric mixing in the absence of phase transitions, and the second stage is the response of the droplets in a vertically moving adiabatic (i.e., closed) parcel. Analytical results in this section are derived under the following assumptions: (1) only liquid exists in the condensed form (no ice); (2) the cloud parcel rises adiabatically; (3) the droplet size distribution is monodisperse; (4) the growth of droplets is due to water vapor condensation; (5) sedimentation and collision-coalescence of droplets are ignored.

2.1 Liquid water mixing ratio in an adiabatic cloud without mixing

For reference, we begin by deriving the change of liquid water mixing ratio in a rising adiabatic cloud parcel without mixing. Considering a warm cloud parcel with monodisperse cloud droplets rising adiabatically with a constant updraft velocity, the supersaturation development equation is (Lamb and Verlinde, 2011, p. 417)

$$\frac{ds}{dt} = Q_1 w - Q_2 \frac{dq_l}{dt}, \quad (1)$$

where s is supersaturation, w is updraft velocity, and q_l is the liquid water mixing ratio (g kg^{-1}). Q_1 and Q_2 depend on temperature, pressure and other constants (all symbols and expressions are given in the Appendix). The first term on the right side represents the production of supersaturation due to adiabatic cooling due to vertical displacement, while the second term accounts for the supersaturation depletion due to vapor condensation. For monodisperse cloud droplets $q_l = (4/3)\pi\rho_w r_d^3 n_d$, where r_d is the radius of cloud droplet and n_d is number concentration in units of kg^{-1} . Thus, $dq_l/dt = 4\pi\rho_w n_d r_d^2 dr_d/dt = 4\pi\rho_w n_d r_d G s$. Here we use the linear growth for an individual droplet: $r_d dr_d/dt = G s$, where G is the condensation growth parameter (see Appendix).

When supersaturation transients are negligible, e.g., after droplet activation, Eq. (1) leads to linear growth rate of q_l ,

$$\frac{dq_l}{dt} = C_1 w, \quad (2)$$

where $C_1 = Q_1/Q_2$ with the units of m^{-1} (see Appendix). This is the quasi-steady limit, in which the supersaturation is

$$s_{qs} = \frac{Aw}{r_d n_d}, \quad (3)$$

where A is a parameter depending on G , Q_1 and Q_2 (see Appendix). If we assume C_1 is a constant, then q_l can be derived by integration of Eq. (2),

$$q_l = C_1 z + q_{l,i}, \quad (4)$$

where $q_{l,i}$ is the initial liquid water mixing ratio, and $z = \int w dt$ is the displacement of the cloud parcel away from its initial location. The liquid water mixing ratio increases linearly with height and does not depend on the updraft velocity. It should be mentioned that Eq. (4) describes q_l under thermodynamic equilibrium conditions. In reality, a cloud system needs some time (phase relaxation time) to reach the equilibrium state; For liquid clouds the phase relaxation time is usually smaller than 10 s (Korolev and Mazin, 2003).

During the adiabatic process, two physical properties of the cloud parcel will be conserved: total water mass mixing ratio and liquid water potential temperature (Kumar et al., 2014), such that

$$q_{l,i} + q_{v,i} = q_{l,f} + q_{v,f} \quad (5)$$

and

$$T_i - \frac{l_w}{c_p} q_{l,i} = T_f - \frac{l_w}{c_p} q_{l,f}, \quad (6)$$

where q is the water mass mixing ratio (g g^{-1}), T is temperature (K), l_w is the latent heat of vaporization of liquid water (J kg^{-1}) and c_p is the specific heat of air at constant pressure ($\text{J kg}^{-1} \text{K}^{-1}$). Subscripts l and v represent liquid and water vapor, respectively, while subscripts i and f denote the initial and final states of the cloud parcel. We note that, for simplicity, the linearized form of the liquid water potential temperature has been used in Eq. (6); i.e., the pressure dependence has been neglected, which is valid if the cloud thickness is not too large.

2.2 Liquid water mixing ratio in an adiabatic cloud after mixing

Now we consider the mixing of a cloud with dry and clean (aerosol-free) environmental air and subsequent evolution for a closed, rising parcel. We define the mixing fraction as χ , such that χ fraction of cloud air is mixed with $(1-\chi)$ fraction of environmental air. We assume the mixing process is isobaric, and that the timescale for the mixing is much smaller than the timescale for the response of the cloud droplets during the mixing (i.e., homogeneous mixing limit). Therefore after isobaric mixing but before any phase changes, the liquid water mixing ratio should be $\chi q_{l,i}$ and the water vapor mixing ratio should be $\chi q_{v,i} + (1-\chi)q_{v,e}$ and the temperature of the mixed parcel should be $\chi T_i + (1-\chi)T_e$. Subscript e denotes the state of the environmental air. After the mixing, we assume the mixed parcel rises adiabatically again with a constant updraft velocity w_m . For the purposes of this derivation w_m is prescribed and we do not consider the actual buoyancy of the mixed parcel with respect to the environment. Similar to Eqs. (5) and (6), we have two conservation equations that allow the liquid water mixing ratio and temperature to be determined for the final state of the mixed parcel (Kumar et al., 2014), denoted by subscript fm:

$$\chi (q_{l,i} + q_{v,i}) + (1-\chi) q_{v,e} = q_{l,fm} + q_{v,fm} \quad (7)$$

and

$$\chi T_i + (1-\chi) T_e - \frac{l_w \chi}{c_p} q_{l,i} = T_{fm} - \frac{l_w}{c_p} q_{l,fm}. \quad (8)$$

Now we ask the following: how does the liquid water mixing ratio in the mixed parcel ($q_{l,fm}$) change with height above the mixing level? What is the difference of liquid water mixing ratio in the mixed parcel ($q_{l,fm}$) compared with that in the original parcel without mixing ($q_{l,f}$) at the same height? How does the difference ($q_{l,f} - q_{l,fm}$) change with height? To calculate this difference, we first subtract Eq. (7) from Eq. (5) to get the liquid water difference for the final state,

$$q_{l,f} - q_{l,fm} = (1-\chi) (q_{l,i} + q_{v,i} - q_{v,e}) - (q_{v,f} - q_{v,fm}). \quad (9)$$

The first term on the right side is the total water mixing ratio difference between the original and new parcel, which depends on the initial condition of the parcel ($q_{l,i}$, $q_{v,i}$), the environmental air ($q_{v,e}$), and the mixing fraction χ . This difference is large when χ is small and environmental air is dry. The second term on the right side is the water vapor mixing ratio difference. The water vapor mixing ratio can be calculated from the temperature, pressure and saturation ratio: $q_v = S e_s(T) \epsilon / (p - e_s(T))$. Therefore the difference of water vapor mixing ratio is

$$q_{v,f} - q_{v,fm} = \frac{S_f e_s(T_f) \epsilon}{p_f - e_s(T_f)} - \frac{S_{fm} e_s(T_{fm}) \epsilon}{p_{fm} - e_s(T_{fm})}. \quad (10)$$

This equation is accurate but not simple enough to be useful. To achieve an analytical result, we first assume $p_f \approx p_{fm}$ because both parcels are at the same height. Secondly, we ignore e_s in the denominator because $p \gg e_s$. In addition, we assume both parcels are in quasi-steady state at that level and that the quasi-stationary supersaturation is much smaller than 1, so that the influence of the change of s_{qs} can be ignored compared with the change of $e_s(T)$ due to temperature; thus we assume $S_{fm} \approx S_f$. The main difference in q_v arises from $e_s(T)$ due to the temperature difference. Using the linearized form of the Clausius–Clapeyron equation, we can approximate the difference of $e_s(T)$ as

$$e_s(T_f) - e_s(T_{fm}) \approx \frac{e_s(T_f) l_w}{R_v T_f^2} (T_f - T_{fm}). \quad (11)$$

From the above assumptions and Eq. (11), we can simplify Eq. (10),

$$q_{v,f} - q_{v,fm} \approx \frac{S_f e_s(T_f) l_w \epsilon}{p_f R_v T_f^2} (T_f - T_{fm}). \quad (12)$$

Combining Eqs. (9) and (12), we find that the liquid water mixing ratio difference depends on the temperature difference in this way,

$$q_{l,f} - q_{l,fm} = (1 - \chi) (q_{l,i} + q_{v,i} - q_{v,e}) - \frac{S_f e_s(T_f) l_w \epsilon}{p_f R_v T_f^2} (T_f - T_{fm}). \quad (13)$$

In addition, the difference in liquid water potential temperature conservation equations for closed and mixed parcels, given by Eq. (6) minus Eq. (8), leads to

$$(1 - \chi) \left(T_i - T_e - \frac{l_w}{c_p} q_{l,i} \right) = T_f - T_{fm} - \frac{l_w}{c_p} (q_{l,f} - q_{l,fm}). \quad (14)$$

Finally, from Eqs. (13) and (14), we can obtain the approximate solutions for liquid water mixing ratio difference and

temperature difference,

$$q_{l,f} - q_{l,fm} = (1 - \chi) \frac{(1 + C_3) q_{l,i} + q_{v,i} - q_{v,e} - C_2 (T_i - T_e)}{1 + C_3} \quad (15)$$

and

$$T_f - T_{fm} = (1 - \chi) \frac{C_2 (T_i - T_e) + C_3 (q_{v,i} - q_{v,e})}{C_2 (1 + C_3)}. \quad (16)$$

Combining Eqs. (4) and (15), we can get the liquid water profile for the mixed parcel,

$$q_{l,fm}(z) = C_1 z + q_{l,i} - (1 - \chi) K_1, \quad (17)$$

where $K_1 = ((1 + C_3) q_{l,i} + q_{v,i} - q_{v,e} - C_2 (T_i - T_e)) / (1 + C_3)$. It is interesting to see that the liquid water mixing ratio for the mixed parcel still increases linearly with height, but with a smaller value compared with an unmixed parcel (cf. Eq. 4). The difference is the same at different heights and is proportional to $1 - \chi$.

2.2.1 Total evaporation and reactivation height

Another way to look at Eq. (17) is that the liquid water mixing ratio in the mixing parcel $q_{l,fm}$ increases with height linearly with the same slope as $q_{l,f}$ in the original parcel, but with a smaller initial liquid water mixing ratio in the mixing parcel $q_{l,im} = q_{l,i} - (1 - \chi) K_1$. Although the initial liquid water mixing ratio $q_{l,im}$ should be non-negative physically, $q_{l,i} - (1 - \chi) K_1$ can be negative mathematically. If $q_{l,im}$ is negative, it means that all cloud droplets evaporate. Therefore, $q_{l,i} = (1 - \chi) K_1$ is the criterion or critical condition that all droplets totally evaporate and the air in mixing parcel is just saturated. This critical condition is consistent with that given by Pinsky et al. (2016b), with $\gamma = 0$.

Even if $q_{l,im}$ is negative at $z = 0$, it can become positive at higher altitude. The negative value of $q_{l,im}$ at the beginning is the result of total evaporation, while the point where $q_{l,im}$ changes to positive can be taken to represent the re-activation of cloud condensation nuclei to form cloud droplets. The re-activation height z_{react} is the distance between the mixing level and the level at which $q_{l,im} = 0$, given by

$$z_{\text{react}} = \frac{(1 - \chi) K_1 - q_{l,i}}{C_1}. \quad (18)$$

2.2.2 Critical height for superadiabatic droplet growth

In this subsection we consider how cloud droplet size changes with height above the mixing level. We consider an initially adiabatic cloud parcel mixed isobarically with clean environmental air at some level above the cloud base. Without vertical movement, the liquid water mixing ratio and cloud number concentration will decrease due to dilution (not considering, for the moment, scenarios in which all droplets

are evaporated). The mean cloud droplet size after the response to mixing is the same for extremely inhomogeneous mixing but smaller for homogeneous mixing. If the parcel still rises adiabatically after mixing, however, the liquid water mixing ratio will increase with height (cf. Eq. 17). This indicates that cloud droplet size will also increase with height, because the number concentration does not change during the vertical motion. We now consider the growth of cloud droplets under quasi-steady conditions. Because the cloud droplet concentration is smaller in the mixed parcel than in the original parcel, s_{qs} in the mixed parcel will be larger ($s_{qs} \propto (r_d n_d)^{-1}$, see Eq. 3). This implies that cloud droplets in the mixed parcel grow faster than those in the original one due to higher supersaturation. Therefore, although cloud droplet size in the mixed parcel is smaller for homogeneous mixing at the beginning, it can, with adequate vertical displacement, become equal to or even larger than that in the original, unmixed parcel. The resulting droplets would appear to have experienced super-adiabatic growth compared to a closed parcel. This general picture of large-drop production resulting from decreased competition in diluted parcels has been discussed elsewhere in the literature (Paluch and Knight, 1986; Su et al., 1998; Cooper et al., 2013; Schmeissner et al., 2015).

We seek a condition to define a critical height z^* at which droplets in the unmixed and mixed parcels have the same radius. Under the assumptions of clean environment and homogeneous mixing, super-adiabatic droplets will exist for $z > z^*$. For the same assumptions and not considering complete evaporation, it is true that $n_{d,fm}/n_{d,f} = \chi$, where χ is the cloud mixing fraction defined in Sect. 2.2. It follows that, for monodisperse clouds, the two parcels will have the same cloud droplet radius when $q_{1,fm}/q_{1,f} = \chi$, because $q_1 \propto r_d^3 n_d$. (Note that this condition for equal radius is the ratio of q_1 , not the difference in q_1 that was shown previously to be constant after mixing, i.e., Eq. 15.) Using Eqs. (17) and (4), the ratio is

$$\frac{C_1 z^* + q_{1,i} - (1 - \chi) K_1}{C_1 z^* + q_{1,i}} = \chi. \quad (19)$$

Solving Eq. (19), we obtain

$$z^* = \frac{K_1 - q_{1,i}}{C_1}. \quad (20)$$

We note with interest that z^* is independent of the mixing fraction χ . Equations (17) and (20) indicate that although the liquid water mixing ratio for the mixed parcel is always lower than that in the original parcel, droplet radius in the mixed parcel will be larger than that in the original parcel when the parcel is above z^* .

3 Results from parcel model

The analytical results derived in Sect. 2 have provided insight into the evolution of a cloud parcel after a mixing event,

but several assumptions and simplifications were made, and some limitations such as perfectly clean (aerosol-free) environment were imposed. We now explore the same concept of idealized mixing and subsequent growth, but using an adiabatic parcel model with bin microphysics. The model was originally developed by Feingold et al. (1998) to simulate warm cloud process and has been applied to a wide range of microphysical problems (Feingold and Kreidenweis, 2000; Xue and Feingold, 2004; Ervens et al., 2005; Ervens and Feingold, 2012; Yang et al., 2012; Li et al., 2013). To study the mixing process, we add an idealized entrainment/detrainment and mixing process to the model. Entrainment means some fraction of environment air flows into the cloud, while detrainment means some fraction of cloud flows into the environment (de Rooy et al., 2013). The mixing process is implemented so that the entrained environmental air is homogeneously mixed with the remaining cloud body, and in all cases considered here this mixing level is set to 665 m (50 m above cloud base). It should be mentioned that mixing process might not necessarily happen when entrainment/detrainment occurs. The time interval between these two processes is called the mixing timescale, and the presence of a delay would be expected for inhomogeneous mixing. The relative magnitudes of this mixing timescale and the phase relaxation time determine whether the mixing occurs in the homogeneous or inhomogeneous limit (e.g., Baker et al., 1980). To be consistent with the previous theoretical development (Sect. 2) we implement the homogeneous mixing limit within the model, i.e., the instantaneous exposure of droplets to the mixture of cloudy and entrained air. This implies that the turbulent mixing time is very small compared to the phase relaxation time and is therefore similar to the limit considered by Pinsky et al. (2016b).

Initial conditions for the parcel are $z_0 = 300$ m, $p_0 = 919$ Pa, $T_0 = 288.15$ K and $RH_0 = 85\%$. CCN are ammonium sulfate particles with a monodisperse radius of 50 nm and concentration of 50 mg^{-1} . The parcel rises adiabatically with constant updraft velocity. Two updraft velocities (w) are chosen in this study: 0.1 and 1.0 m s^{-1} . Observation results show that updraft velocity in cumulus cloud is on the order of 1.0 m s^{-1} , and that for stratocumulus cloud is on the order of 0.1 m s^{-1} (Katzwinkel et al., 2014; Ditas et al., 2012). Cloud base is reached at $z = 615$ m, where CCN are activated as cloud droplets. The isobaric mixing process occurs at $z = 665$ m, 50 m above the cloud base. For simplicity, we assume the environmental temperature at the mixing level is the same as that of the cloud parcel, but the relative humidity is only 85%. After the mixing, the new mixed parcel rises adiabatically again with the same updraft velocity.

Liquid water mixing ratio profiles for six different mixing fractions $\chi = 1.0, 0.9, 0.8, 0.7, 0.6, 0.5$ at $w = 0.1 \text{ m s}^{-1}$ are shown in Fig. 1a. The analytical results based on Eq. (17) are also shown and are quite close to the results from the parcel model. As seen from Fig. 1a, the liquid water mixing ratio for smaller χ is smaller than that for larger χ at the




Figure 1. (a) Liquid water mixing ratio profiles for various cloud mixing fractions χ and with low updraft speed (0.1 m s^{-1}). Lines are from the parcel model and dots are from the theoretical prediction given by Eq. (17). (b) Cloud droplet radius vs. height for various χ when mixing with clean (aerosol-free) environmental air. The horizontal dashed line represents the critical height z^* calculated from Eq. (20). (c) Cloud droplet radius vs. height for various χ when mixing with polluted environmental air (air containing CCN similar to cloud base conditions). Insets in panels (b) and (c) show details of the radius profiles above the critical height. Super-adiabatic droplet growth, i.e., droplet diameters greater than in the unmixed cloud ($\chi = 1.0$), is observed for all χ in panel (b) and only for $\chi = 0.9$ in panel (c).

same height. In addition, when $\chi \leq 0.8$, the liquid water mixing ratio will decrease to zero at the beginning, which means that the cloud totally evaporates and becomes subsaturated. It should be mentioned that in the model each cloud droplet contains one CCN, and when a cloud droplet totally evaporates the CCN still survives. Because the subsaturated parcel still rises adiabatically, CCN in the mixing parcel can be activated again when the air becomes saturated at a higher level, which we defined as the re-activation level. The smaller χ is, the higher the re-activation level is. The evaporation and re-activation processes can be clearly seen from the cloud droplet radius profile in Fig. 1b. In addition, it clearly shows that the mixed cloud parcel can reach super-adiabatic growth conditions (where the cloud droplet radius in the mixed parcel is larger than that in the original, unmixed parcel with $\chi = 1.0$) above a critical height. The critical height is independent of χ and agrees well with that predicted by Eq. (20). The saturation ratio and cloud droplet number concentration profiles for this case are shown in Fig. S1 (in the Supplement). It can be seen that cloud droplet number concentration in the mixed parcel decreases with decreasing χ , while supersaturation increases with decreasing χ in the quasi-steady region.

Results above are for a cloud parcel mixing with clean environmental air (aerosol-free condition). However, both observational and modeling results show that air around the cumulus cloud is usually not clean (Katzwinkel et al., 2014; Chen et al., 2012). There can be background aerosols in the

atmosphere even at high altitude, and in addition subsiding shells can also provide sufficient aerosols as CCN when mixing occurs (Heus and Jonker, 2008). There is no simple analytical result for mixing with a polluted environment. However, we can use the parcel model to investigate the effect of mixing when the environmental air is polluted. For simplicity, we assume the environment has the same dry aerosol size distribution as that below the cloud base.

Figure 1c shows the monodisperse cloud droplet radius vs. height for various χ after mixing with a polluted environment at $w = 0.1 \text{ m s}^{-1}$. For $\chi = 0.9$, the remaining cloud droplets do not totally evaporate and the entrained aerosols are not activated as cloud droplets. For smaller χ , the remaining cloud droplets totally evaporate and leave CCN in the mixed parcel. Both entrained and remaining CCN are activated at a higher level. In addition, only the parcel with $\chi = 0.9$ can reach the super-adiabatic growth region. For smaller χ , cloud droplets are smaller than those in the original parcel at the same height z^* . In summary, when mixing with a polluted environment, the mixing parcel can reach super-adiabatic growth conditions at the predicted z^* only if the cloud does not totally evaporate after mixing. Saturation ratio and cloud droplet concentration profiles for various mixing fractions are shown in Fig. S2. Supersaturation in the quasi-steady state is smaller than that when mixing with the clean environmental air as shown in Fig. S1. This is because the entrained aerosols from the polluted environmental air can be activated as cloud




Figure 2. (a) Liquid water mixing ratio profiles for various cloud mixing fractions χ and with high updraft speed (1.0 m s^{-1}). Lines are from the parcel model and dots are from the theoretical prediction given by Eq. (17). (b) Cloud droplet radius vs. height for various χ when mixing with clean (aerosol-free) environmental air. The horizontal dashed line represents the critical height z^* calculated from Eq. (20). (c) Cloud droplet radius vs. height for various χ when mixing with polluted environmental air (air containing CCN similar to cloud base conditions). Insets in panels (b) and (c) show details of the radius profiles above the critical height. Super-adiabatic droplet growth, i.e., droplet diameters greater than in the unmixed cloud ($\chi = 1.0$), is observed for all χ in panel (b) but for none in panel (c).

droplets and thus suppress the supersaturation in the mixed parcel.

Figure 2a and b show the results for mixing with a clean environment at larger updraft velocity $w = 1.0 \text{ m s}^{-1}$. It can be seen that the liquid water mixing ratio and cloud droplet radius profiles are almost the same compared with Fig. 1, except that the mixing parcel totally evaporate for $\chi = 0.8$ at $w = 0.1 \text{ m s}^{-1}$, but does not totally evaporate for $\chi = 0.8$ at $w = 1.0 \text{ m s}^{-1}$. This is because larger updraft velocity supplies more water within the fixed phase relaxation time, so droplets begin to grow before they have had time to completely evaporate. The mixed parcel can reach the super-adiabatic growth region when it is above z^* . And as before, z^* is independent of both mixing fraction and updraft velocity, consistent with the theoretical prediction.

When mixing with polluted environment air at $w = 1.0 \text{ m s}^{-1}$, the mixed parcel cannot reach the super-adiabatic growth region, whether the mixing parcel totally evaporates or not (see Fig. 2c). The reason is that with large updraft velocity, the entrained CCN can always be activated as cloud droplets, thus competing for water vapor in the mixed parcel. It should be mentioned that results here strongly depend on the physical and chemical properties of the entrained CCN, e.g., sizes, chemical composition, and number concentration. For example, the mixed parcel might also reach the super-adiabatic growth region if the environmental air only contains a small number of CCN. In general, however, mixing with polluted air will inhibit the super-adiabatic growth

of cloud droplets. Saturation ratio and cloud droplet number concentration profiles for clean and polluted conditions with high updraft velocity are shown in Figs. S3 and S4 separately.

Cloud droplets in a real cloud are usually polydisperse and we now consider to what extent the theoretical predictions apply in this more complex system. The effect of mixing on a polydisperse droplet population is tested with the cloud parcel model. The initial aerosols are composed of ammonium sulfate and are distributed lognormally in 20 bins with 50 nm median radius, standard deviation of 1.4, and a total number concentration of 100 cm^{-3} . Initial radii of the dry aerosols for the 20 bins are listed in the Supplement. The cloud droplet diameters for each bin vs. height for $\chi = 0.9, 0.7, 0.5$ are shown in Fig. 3. These results are for clean environmental air and $w = 0.1 \text{ m s}^{-1}$ and are representative of the other cases. It can be seen that not all 20 bins are activated at cloud base; for example, only the largest 11 aerosol sizes are activated as cloud droplets for $\chi = 1.0$. Cloud droplets evaporate a little bit for $\chi = 0.9$, or completely for $\chi = 0.7, 0.5$, and reactivation occurs again at a higher level. It is very interesting to see that for $\chi = 0.5$, the 12th bin is not activated at cloud base but is activated for the first time after mixing (green line). This asymmetric phenomenon is due to the significant reduction of cloud droplet number concentration after mixing. Thermodynamic equilibrium predicts how much water vapor should condense at a certain level, but mixing with a clean environment reduces the overall CCN concentration. To condense the same amount of water, either the single




Figure 3. Radii of cloud droplets in a polydisperse population vs. height for $\chi = 0.9, 0.7, 0.5$ in a clean environment at $w = 0.1 \text{ m s}^{-1}$. The background grey lines represent $\chi = 1.0$. The right column shows the region near the critical height where super-adiabatic growth can be expected. The black line is for the 1st bin (largest CCN), the red line for the 11th bin, and the green line for the 12th bin.

droplets must grow larger than before, which is the physical explanation for super-adiabatic growth; or some initially un-activated aerosol particles can be activated to increase the cloud number concentration.

Super-adiabatic droplet growth for individual droplet size bins can be observed in Fig. 3, but it is achieved at different heights above the mixing level. Figure 4 shows these critical heights for individual cloud droplet size bins calculated from the cloud parcel model for the various mixing fractions. Here again, the environmental air is clean with $T_e = T_c$ and $\text{RH}_e = 85\%$. We note that cloud droplet size decreases with increasing bin number (i.e., cloud droplet size increases with increasing dry aerosol size, as expected). The critical height for each bin is defined when the sizes of cloud droplets for that bin are equal for both mixed and unmixed cloud parcels. It can be seen that the critical height depends on the size of the cloud droplet, the mixing fraction and the updraft velocity, especially for low updraft velocity $w = 0.1 \text{ m s}^{-1}$. For $w = 1.0 \text{ m s}^{-1}$, critical heights for individual bins are close to the theoretical critical height for a monodisperse cloud droplet population. In the low updraft speed case (left panel) it is particularly striking that the $\chi = 0.9$ curve has a different dependence than that for the other mixing fractions: increasing rather than decreasing z^* with decreasing droplet size. We believe the explanation is that the $\chi = 0.9$ case is the only scenario in which complete droplet evaporation does not occur. Thus, the presence of complete evaporation and




Figure 4. Critical height for individual droplet size bins for a polydisperse cloud droplet population calculated from cloud parcel model. Results are shown for two updraft velocities, (a) $w = 0.1 \text{ m s}^{-1}$ and (b) $w = 1.0 \text{ m s}^{-1}$. The line colors represent different mixing fractions χ as defined in the legend, and the dashed line is the analytical result for critical height z^* for a monodisperse cloud droplet population. Cloud droplet size decreases as the bin number increases.

subsequent re-activation changes the population dynamics of the cloud substantially for low updraft speeds. Although the critical heights are different for individual size bins, we might expect that the simple monodisperse prediction for z^* would hold for some moment of cloud droplet size distribution. Considering that the thermodynamically predicted water mass is distributed over a variable number of aerosol particles, the most logical choice is a prediction of z^* using the volume-mean radius. Figure 5 shows the volume-weighted mean radius as a function of height for six values of χ and for updraft speeds of 0.1 and 1.0 m s^{-1} . In spite of the complex behavior observed for individual bins, the volume-mean radius curves are observed to cross at nearly the same height and with very close agreement with the analytical prediction. This suggests that the theory can be applied under realistic cloud conditions with polydisperse droplet populations. Figures S5 and S6 show the saturation ratio and cloud droplet number concentration profiles for polydisperse cloud droplets at low and high updraft velocity separately. Our results are similar to Wang et al. (2009), where they observed faster droplet growth resulting from reduced droplet number concentration and increased supersaturation in a mixed parcel.

4 Discussion and conclusions

In this study, we have considered isobaric mixing of a cloud parcel with environmental air, and then the subsequent droplet growth as the parcel rises adiabatically afterwards.




Figure 5. Volume-mean radius for a polydisperse cloud droplet population vs. height at updraft speeds of (a) $w = 0.1 \text{ m s}^{-1}$ and (b) $w = 1.0 \text{ m s}^{-1}$ and for a clean environment. Line colors represent different mixing fractions χ , as in Figs. 1 and 2. The horizontal dashed line is the critical height z^* predicted for a monodisperse cloud droplet population with equal volume-mean radius.

Analytical expressions are derived for monodisperse cloud droplets when mixing with clean environmental air. Results show that the liquid water mixing ratio q_l in the mixed parcel increases linearly with height with the same slope (dq_l/dz) as the original parcel (without mixing). Due to the mixing the q_l is smaller compared with the unmixed parcel at the same height. A closed form expression for the offset is derived and shows that the decrease of q_l in the mixed parcel depends on the mixing fraction χ and the temperature and relative humidity of the environmental air. A critical height z^* , defined as the height at which the cloud droplet sizes are equal in both mixed and original cloud parcels, is derived. Interestingly, the critical height depends on the initial conditions of the cloud and environmental air, but is independent of the mixing fraction. Cloud droplets in the mixed parcel are larger than in the original parcel above z^* , which we call the “super-adiabatic” growth region. These large cloud droplets may help explain the formation of initial large droplets that contribute to precipitation formation in warm clouds.

The predicted vertical profile of liquid water mixing ratio and the critical height are confirmed using a bin microphysical cloud model. For large χ and a humid environment, cloud droplets will evaporate a little bit and grow again after mixing. For small χ and dry environment, cloud droplets can evaporate completely, leaving the mixed parcel subsaturated. Droplets are re-activated at a higher level, as long as the mixing parcel rises sufficiently to reach saturation again. The theoretical predictions are based on equilibrium arguments, but because the phase relaxation time is typically short for warm clouds, results are not very sensitive to updraft speed over the range investigated. For monodisperse cloud droplets, z^* is

independent of mixing fraction and updraft speed. For polydisperse cloud droplets, however, z^* defined for individual droplet sizes is observed to depend on droplet size, mixing fraction and updraft velocity, especially for smaller w . For larger w , z^* is insensitive to those variables and close to the analytical result for monodisperse cloud droplets. The situation becomes much simpler and the polydisperse cloud can be predicted theoretically by using the volume-mean cloud droplet radius. Finally, we note that the model results presented here are for the condition of cloud and environment having the same temperature; model runs for other reasonable conditions also produced good agreement with the theory.

Environment background aerosols and subsiding shells may contain effective CCN that can be activated after mixing, thus inhibiting super-adiabatic droplet growth. For large updraft speed, the entrained aerosols can be activated as cloud droplets, thus increasing cloud droplet concentration and decreasing the cloud droplet sizes. For small updraft velocity, the mixed parcel can reach the super-adiabatic growth region only when the entrained aerosols cannot be activated and the cloud droplets do not totally evaporate. Otherwise if cloud droplets totally evaporate, both remaining and entrained CCN can be activated when the mixed parcel is saturated again. If the entrained aerosols can be activated as cloud droplets, the mixed parcel usually contains smaller cloud droplets, but similar number concentration compared with the main cloud body. This might help explain the observation that some cloud samples appear to be undiluted in droplet number concentration, but have significantly smaller mean-volume radii, a region otherwise outside the homogeneous mixing limiting curve in a mixing diagram (Schmeisser et al., 2015).

Given the success of the analytical results in predicting the critical height z^* above which volume-weighted mean droplet diameters will appear to be super-adiabatic, we briefly explore the dependence of z^* on environmental conditions. As noted already, and now confirmed by the parcel model, the critical height does not depend on mixing fraction χ or on the updraft speed w . As seen in Fig. 6, z^* changes with the relative humidity of the environmental air (RH_e) at the mixing level. It can be seen that z^* decreases as RH_e increases. For example, when $\text{RH}_e = 98\%$, z^* is less than 50 m above the mixing level. This means that the mixed parcel can reach the super-adiabatic growth region more easily when mixing with a humid environment. Thus the results are relevant to shallow convective clouds, in contrast with the particular example chosen for Figs. 1–5 that requires a height of approximately 300 m above the mixing level for super-adiabatic growth. In the real atmosphere, this has relevance for at least two scenarios. First, for cumulus convection the subsiding shell can be very humid due to the evaporation of cloud droplets at higher cloud levels (Katzwinkel et al., 2014). Mixing under these conditions would be favorable for super-adiabatic growth of cloud droplets, especially if the




Figure 6. Critical height z^* vs. environmental relative humidity RH_e at the mixing level. The height of cloud base (blue dashed line) and the mixing level (black dashed line) are shown for reference.

subsiding shell has been cleared of most CCN through scavenging. Second, for stratocumulus convection the concepts here can hold for mixing between undiluted cloud parcels and parcels previously diluted through cloud-top mixing (followed by descent together with cloud droplet evaporation and humidification). Upon subsequent lifting after mixing with the diluted but humid parcel, super-adiabatic droplets can be produced. This mechanism has similarities to the cycles of vertical motion and repeated mixing described by Wang et al. (2009) and Korolev et al. (2013).

The results presented here all are for the homogeneous mixing limit. It is possible to develop model prescriptions for extreme inhomogeneous mixing, but our sense is that the results would be sensitive to the necessarily artificial nature of those prescriptions. Ultimately, a realistic model or a direct numerical simulation of the mixing process are required for the inhomogeneous limit. We can speculate, however, that the results obtained here would only be amplified for inhomogeneous mixing: in that limit the droplet concentration is reduced but the mean volume diameter remains unchanged, implying that z^* is zero and super-adiabatic droplet growth can begin immediately after the mixing process has concluded. By concluded we mean that the cloudy and environmental air have become completely mixed, leaving a spatially homogeneous field of droplets having the same diameter as before mixing, but lower number concentration due to dilution and total evaporation of some subset of droplets (e.g., Beals et al., 2015). This neglects the more complicated interactions that might come into play if CCN are entrained during mixing with environmental air: in that case activation of new CCN may occur as the parcel rises, even before complete mixing to the microscale has taken place.

A crucial factor that has not been considered thus far is the influence of mixing on the vertical motion of a cloud parcel

due to changes in its buoyancy. Whether a mixed cloud parcel can experience super-adiabatic droplet growth depends not only on the critical height z^* but also on the maximum height z_{\max} it can reach after mixing: a cloud can reach the super-adiabatic growth region only for $z_{\max} > z^*$. Calculation of z_{\max} is nontrivial because one must consider the time dependence of the buoyancy, drag force, and kinetic energy of the parcel, which depends on the properties of the surrounding environment and its dependence on height. These are still open research problems (e.g., slippery vs. sticky thermals (Sherwood et al., 2013; Romps and Charn, 2015)), so exploring this important aspect is beyond the scope of our paper; but qualitatively, our results imply that strongly convective clouds may favor super-adiabatic growth compared to weakly convective clouds. In addition, decreasing χ will tend to decrease the buoyancy and therefore the updraft speed, thus ultimately decreasing z_{\max} . Therefore, it is more likely to reach the super-adiabatic droplet growth region for larger χ , again favoring clouds in humid environments or clouds with well developed, humid subsiding shells.

In a real cloud the liquid water mixing ratio profile is much more complicated than considered here. Mixing will occur at different levels and environmental conditions change with height. There are several methods to predict the mixing fraction at different levels. For example, Lu et al. (2012) predict χ using the cloud base condition, liquid water mixing ratio and environmental condition at each level. The advantage of their method is that they do not need to measure temperature and water vapor mixing ratio in the cloud, which have significant measurement uncertainty. Here, we have provided an explicit method to estimate the mixing fraction at each level using a similar strategy. Based on Eqs. (15) and (16), we can also calculate the mixing fraction profile. The key difference is that our method is explicit, while their method is implicit.

The central insights of this work are the derived critical height z^* for super-adiabatic growth (Eq. 20) and the findings that a mixed parcel is more likely to reach the super-adiabatic growth region when convection is strong, and the environmental air is humid and clean. Cloud droplets in the super-adiabatic growth region $z > z^*$ are larger than that in an unmixed parcel. The theoretical results obtained here and confirmed with the parcel model are a step toward evaluating the possible role of mixing-induced droplet growth for large droplet production and development of precipitation in warm clouds.

5 Data availability

The main results of this paper can be obtained from the derived equations. Results of the parcel model runs described in this work are available upon request from Fan Yang (fyang6@mtu.edu).

Appendix A: List of symbols

Symbol	Description	Units
A	$\frac{Q_1}{4\pi\rho_w G Q_2}$	s kg^{-1}
c_p	specific heat of air at constant temperature	$\text{J kg}^{-1} \text{K}^{-1}$
C_1	$4\pi\rho_w G A = Q_1/Q_2$	m^{-1}
C_2	$\frac{S_f e_s(T_i) l_w \epsilon}{p_f R_v T_i^2}$	K^{-1}
C_3	$\frac{C_2 l_w}{c_p}$	–
D_v	diffusivity of water vapor	$\text{m}^2 \text{s}^{-1}$
e_v	water vapor pressure	Pa
$e_s(T)$	saturated water vapor pressure at temperature T	Pa
G	$\left[\frac{\rho_w R_v T}{D_v e_s(T)} + \frac{\rho_w l_w}{k_T T} \left(\frac{l_w}{R_v T} - 1 \right) S \right]^{-1}$	$\text{m}^2 \text{s}^{-1}$
k_T	coefficient of air heat conductivity	$\text{J m}^{-1} \text{s}^{-1}$
K_1	$\frac{(1+C_3)q_{1,i}+q_{v,i}-q_{v,e}-C_2(T_i-T_e)}{1+C_3}$	–
K_2	$\frac{C_2(T_i-T_e)+C_3(q_{v,i}-q_{v,e})}{C_2(1+C_3)}$	K
l_w	latent heat of liquid water	J kg^{-1}
M_{air}	molar mass of air	kg mol^{-1}
M_w	molar mass of water	kg mol^{-1}
n_d	droplet number per unit mass of air	kg^{-1}
q_1	liquid water mixing ratio	–
$q_{1,i}$	initial q_1	–
$q_{1,f}$	final q_1	–
q_v	water vapor mixing ratio	–
$q_{v,e}$	environmental q_v	–
$q_{v,i}$	initial q_v	–
$q_{v,f}$	final q_v	–
Q_1	$\frac{g l_w}{c_p R_v T^2} - \frac{g}{R_a T}$	m^{-1}
Q_2	$\frac{\rho_{\text{air}} l_w^2}{p c_p T} + \frac{\rho_{\text{air}} R_v T}{e_s(T)}$	–
r_d	radius of cloud droplet	m
$r_{d,i}$	initial r_d	m
$r_{d,f}$	final r_d	m
$r_{d,\text{fm}}$	final r_d with mixing fraction χ	m
R	universal gas constant	$\text{J mol}^{-1} \text{K}^{-1}$
R_a	gas constant for dry air	$\text{J kg}^{-1} \text{K}^{-1}$
R_v	gas constant for water vapor	$\text{J kg}^{-1} \text{K}^{-1}$
s	$S - 1$, water vapor supersaturation	–
S	$\frac{e_v}{e_s}$, water vapor saturation ratio	–
S_f	final S	–
S_{fm}	final S with mixing fraction χ	–
T	temperature	K
T_i	initial T	K
T_{im}	initial T with mixing fraction χ	K
T_e	environmental T	K
T_f	final T	K
T_{fm}	final T with mixing fraction χ	K
w	updraft velocity of cloud parcel	m s^{-1}
w_m	updraft velocity of cloud parcel with mixing fraction χ	m s^{-1}
χ	isobaric mixing fraction	–
ϵ	$\frac{M_w}{M_{\text{air}}}$	–
κ	$\frac{R}{c_p}$	–
ρ_w	density of liquid water	kg m^{-3}
ρ_{air}	density of air	kg m^{-3}

The Supplement related to this article is available online at doi:10.5194/acp-16-9421-2016-supplement.

Acknowledgements. This research was supported by the DOE Office of Science as part of the Atmospheric System Research program through grant no. DE-SC0011690.

Edited by: T. Garrett

Reviewed by: two anonymous referees

References

- Andrejczuk, M., Grabowski, W. W., Malinowski, S. P., and Smolarkiewicz, P. K.: Numerical simulation of cloud-clear air interfacial mixing: Effects on cloud microphysics, *J. Atmos. Sci.*, 63, 3204–3225, 2006.
- Andrejczuk, M., Grabowski, W. W., Malinowski, S. P., and Smolarkiewicz, P. K.: Numerical simulation of cloud-clear air interfacial mixing: homogeneous vs. inhomogeneous mixing, *J. Atmos. Sci.*, 66, 2493–2500, 2009.
- Baker, M., Corbin, R., and Latham, J.: The influence of entrainment on the evolution of cloud droplet spectra: I. A model of inhomogeneous mixing, *Q. J. Roy. Meteor. Soc.*, 106, 581–598, 1980.
- Beals, M. J., Fugal, J. P., Shaw, R. A., Lu, J., Spuler, S. M., and Stith, J. L.: Holographic measurements of inhomogeneous cloud mixing at the centimeter scale, *Science*, 350, 87–90, 2015.
- Beard, K. V. and Ochs III, H. T.: Warm-rain initiation: An overview of microphysical mechanisms, *J. Appl. Meteorol.*, 32, 608–625, 1993.
- Blyth, A. M., Lasher-Trapp, S. G., Cooper, W. A., Knight, C. A., and Latham, J.: The role of giant and ultragiant nuclei in the formation of early radar echoes in warm cumulus clouds, *J. Atmos. Sci.*, 60, 2557–2572, 2003.
- Burnet, F. and Brenguier, J.-L.: Observational study of the entrainment-mixing process in warm convective clouds, *J. Atmos. Sci.*, 64, 1995–2011, 2007.
- Chen, G., Xue, H., Feingold, G., and Zhou, X.: Vertical transport of pollutants by shallow cumuli from large eddy simulations, *Atmos. Chem. Phys.*, 12, 11319–11327, doi:10.5194/acp-12-11319-2012, 2012.
- Chen, J., Liu, Y., and Zhang, M.: Investigation the influences of entrainment mixing processes on cloud microphysics using new cloud parcel model, AGU Fall Meeting, 18 December 2015, San Francisco, USA, A51D-0083, 2015.
- Cheng, W. Y., Carrió, G. G., Cotton, W. R., and Saleeby, S. M.: Influence of cloud condensation and giant cloud condensation nuclei on the development of precipitating trade wind cumuli in a large eddy simulation, *J. Geophys. Res.-Atmos.* (1984–2012), 114, D08201, doi:10.1029/2008JD011011, 2009.
- Cooper, W. A., Lasher-Trapp, S. G., and Blyth, A. M.: The influence of entrainment and mixing on the initial formation of rain in a warm cumulus cloud, *J. Atmos. Sci.*, 70, 1727–1743, 2013.
- Cotton, W. R., Bryan, G., and Van den Heever, S. C.: Storm and cloud dynamics, vol. 99, Academic press, Burlington, MA, USA, 2011.
- de Lozar, A. and Muessle, L.: Long-resident droplets at the stratocumulus top, *Atmos. Chem. Phys.*, 16, 6563–6576, doi:10.5194/acp-16-6563-2016, 2016.
- de Rooy, W. C., Bechtold, P., Fröhlich, K., Hohenegger, C., Jonker, H., Mironov, D., Pier Siebesma, A., Teixeira, J., and Yano, J.-I.: Entrainment and detrainment in cumulus convection: an overview, *Q. J. Roy. Meteor. Soc.*, 139, 1–19, 2013.
- Ditas, F., Shaw, R. A., Siebert, H., Simmel, M., Wehner, B., and Wiedensohler, A.: Aerosols-cloud microphysics-thermodynamics-turbulence: evaluating supersaturation in a marine stratocumulus cloud, *Atmos. Chem. Phys.*, 12, 2459–2468, doi:10.5194/acp-12-2459-2012, 2012.
- Ervens, B. and Feingold, G.: On the representation of immersion and condensation freezing in cloud models using different nucleation schemes, *Atmos. Chem. Phys.*, 12, 5807–5826, doi:10.5194/acp-12-5807-2012, 2012.
- Ervens, B., Feingold, G., and Kreidenweis, S. M.: Influence of water-soluble organic carbon on cloud drop number concentration, *J. Geophys. Res.-Atmos.* (1984–2012), 110, D18211, doi:10.1029/2004JD005634, 2005.
- Feingold, G. and Kreidenweis, S.: Does cloud processing of aerosol enhance droplet concentrations?, *J. Geophys. Res.-Atmos.* (1984–2012), 105, 24351–24361, 2000.
- Feingold, G., Walko, R., Stevens, B., and Cotton, W.: Simulations of marine stratocumulus using a new microphysical parameterization scheme, *Atmos. Res.*, 47, 505–528, 1998.
- Feingold, G., Cotton, W. R., Kreidenweis, S. M., and Davis, J. T.: The impact of giant cloud condensation nuclei on drizzle formation in stratocumulus: Implications for cloud radiative properties, *J. Atmos. Sci.*, 56, 4100–4117, 1999.
- Gerber, H., Frick, G., Jensen, J., and Hudson, J.: Entrainment, mixing, and microphysics in trade-wind cumulus, *J. Meteor. Soc. Japan*, 86, 87–106, 2008.
- Göke, S., Ochs III, H. T., and Rauber, R. M.: Radar analysis of precipitation initiation in maritime vs. continental clouds near the Florida coast: Inferences concerning the role of CCN and giant nuclei, *J. Atmos. Sci.*, 64, 3695–3707, 2007.
- Heus, T. and Jonker, H. J.: Subsiding shells around shallow cumulus clouds, *J. Atmos. Sci.*, 65, 1003–1018, 2008.
- Hocking, L.: The collision efficiency of small drops, *Q. J. Roy. Meteor. Soc.*, 85, 44–50, 1959.
- Jensen, J., Austin, P., Baker, M., and Blyth, A.: Turbulent mixing, spectral evolution and dynamics in a warm cumulus cloud, *J. Atmos. Sci.*, 42, 173–192, 1985.
- Jensen, J. B. and Lee, S.: Giant sea-salt aerosols and warm rain formation in marine stratocumulus, *J. Atmos. Sci.*, 65, 3678–3694, 2008.
- Johnson, D. B.: The role of giant and ultragiant aerosol particles in warm rain initiation, *J. Atmos. Sci.*, 39, 448–460, 1982.
- Katzwinkel, J., Siebert, H., Heus, T., and Shaw, R. A.: Measurements of turbulent mixing and subsiding shells in trade wind cumuli, *J. Atmos. Sci.*, 71, 2810–2822, 2014.
- Korolev, A., Pinsky, M., and Khain, A.: A New Mechanism of Droplet Size Distribution Broadening during Diffusional Growth, *J. Atmos. Sci.*, 70, 2051–2071, 2013.
- Korolev, A., Khain, A., Pinsky, M., and French, J.: Theoretical study of mixing in liquid clouds – Part 1: Classical concepts, *Atmos. Chem. Phys.*, 16, 9235–9254, doi:10.5194/acp-16-9235-2016, 2016.

- Korolev, A. V. and Mazin, I. P.: Supersaturation of water vapor in clouds, *J. Atmos. Sci.*, 60, 2957–2974, 2003.
- Kostinski, A. B. and Shaw, R. A.: Fluctuations and luck in droplet growth by coalescence, *B. Am. Meteorol. Soc.*, 86, 235–244, 2005.
- Kumar, B., Schumacher, J., and Shaw, R. A.: Lagrangian mixing dynamics at the cloudy-clear air interface, *J. Atmos. Sci.*, 71, 2564–2580, doi:10.1175/JAS-D-13-0294.1, 2014.
- Laird, N. F., Ochs Iii, H. T., Rauber, R. M., and Miller, L. J.: Initial precipitation formation in warm Florida cumulus, *J. Atmos. Sci.*, 57, 3740–3751, 2000.
- Lamb, D. and Verlinde, J.: *Physics and chemistry of clouds*, Cambridge University Press, New York, NY, USA, 2011.
- Lasher-Trapp, S. G., Cooper, W. A., and Blyth, A. M.: Broadening of droplet size distributions from entrainment and mixing in a cumulus cloud, *Q. J. Roy. Meteor. Soc.*, 131, 195–220, 2005.
- Lehmann, K., Siebert, H., and Shaw, R. A.: Homogeneous and inhomogeneous mixing in cumulus clouds: dependence on local turbulence structure, *J. Atmos. Sci.*, 66, 3641–3659, 2009.
- Li, Z., Xue, H., and Yang, F.: A modeling study of ice formation affected by aerosols, *J. Geophys. Res.-Atmos.*, 118, 11–213, 2013.
- Lu, C., Liu, Y., Yum, S. S., Niu, S., and Endo, S.: A new approach for estimating entrainment rate in cumulus clouds, *Geophys. Res. Lett.*, 39, L04802, doi:10.1029/2011GL050546, 2012.
- Lu, C., Niu, S., Liu, Y., and Vogelmann, A. M.: Empirical relationship between entrainment rate and microphysics in cumulus clouds, *Geophys. Res. Lett.*, 40, 2333–2338, 2013.
- Magaritz-Ronen, L., Pinsky, M., and Khain, A.: Effects of Turbulent Mixing on the Structure and Macroscopic Properties of Stratocumulus Clouds Demonstrated by a Lagrangian Trajectory Model, *J. Atmos. Sci.*, 71, 1843–1862, 2014.
- Magaritz-Ronen, L., Pinsky, M., and Khain, A.: Drizzle formation in stratocumulus clouds: effects of turbulent mixing, *Atmos. Chem. Phys.*, 16, 1849–1862, doi:10.5194/acp-16-1849-2016, 2016.
- Naumann, A. K. and Seifert, A.: A Lagrangian drop model to study warm rain microphysical processes in a shallow cumulus, *J. Adv. Model. Earth Syst.*, 7, 1136–1154, 2015.
- Paluch, I. R. and Knight, C. A.: Does mixing promote cloud droplet growth?, *J. Atmos. Sci.*, 43, 1994–1998, 1986.
- Pinsky, M., Khain, A., and Korolev, A.: Theoretical analysis of mixing in liquid clouds – Part 3: Inhomogeneous mixing, *Atmos. Chem. Phys.*, 16, 9273–9297, doi:10.5194/acp-16-9273-2016, 2016a.
- Pinsky, M., Khain, A., Korolev, A., and Magaritz-Ronen, L.: Theoretical investigation of mixing in warm clouds – Part 2: Homogeneous mixing, *Atmos. Chem. Phys.*, 16, 9255–9272, doi:10.5194/acp-16-9255-2016, 2016b.
- Pruppacher, H. R., Klett, J. D., and Wang, P. K.: *Microphysics of clouds and precipitation*, Taylor & Francis, Dordrecht, the Netherlands, 1998.
- Romps, D. M. and Charn, A. B.: Sticky thermals: Evidence for a dominant balance between buoyancy and drag in cloud updrafts, *J. Atmos. Sci.*, 72, 2890–2901, 2015.
- Schmeissner, T., Shaw, R., Ditas, J., Stratmann, F., Wendisch, M., and Siebert, H.: Turbulent Mixing in Shallow Trade Wind Cumuli: Dependence on Cloud Life Cycle, *J. Atmos. Sci.*, 72, 1447–1465, 2015.
- Sherwood, S. C., Hernández-Deckers, D., Colin, M., and Robinson, F.: Slippery Thermals and the Cumulus Entrainment Paradox, *J. Atmos. Sci.*, 70, 2426–2442, 2013.
- Su, C.-W., Krueger, S. K., McMurtry, P. A., and Austin, P. H.: Linear eddy modeling of droplet spectral evolution during entrainment and mixing in cumulus clouds, *Atmos. Res.*, 47, 41–58, 1998.
- Tölle, M. H. and Krueger, S. K.: Effects of entrainment and mixing on droplet size distributions in warm cumulus clouds, *J. Adv. Model. Earth Syst.*, 6, 281–299, 2014.
- Wang, J., Daum, P. H., Yum, S. S., Liu, Y., Senum, G. I., Lu, M.-L., Seinfeld, J. H., and Jonsson, H.: Observations of marine stratocumulus microphysics and implications for processes controlling droplet spectra: Results from the Marine Stratus/Stratocumulus Experiment, *J. Geophys. Res.-Atmos.*, 114, D18210, doi:10.1029/2008JD011035, 2009.
- Xue, H. and Feingold, G.: A modeling study of the effect of nitric acid on cloud properties, *J. Geophys. Res.-Atmos.* (1984–2012), 109, D18204, doi:10.1029/2004JD004750, 2004.
- Yang, F., Xue, H., Deng, Z., Zhao, C., and Zhang, Q.: A closure study of cloud condensation nuclei in the North China Plain using droplet kinetic condensational growth model, *Atmos. Chem. Phys.*, 12, 5399–5411, doi:10.5194/acp-12-5399-2012, 2012.
- Yin, Y., Levin, Z., Reisin, T. G., and Tzivion, S.: The effects of giant cloud condensation nuclei on the development of precipitation in convective clouds – A numerical study, *Atmos. Res.*, 53, 91–116, 2000.
- Yum, S. S., Wang, J., Liu, Y., Senum, G., Springston, S., McGraw, R., and Yeom, J. M.: Cloud microphysical relationships and their implication on entrainment and mixing mechanism for the stratocumulus clouds measured during the VOCALS project, *J. Geophys. Res.-Atmos.*, 120, 5047–5069, doi:10.1002/2014JD022802, 2015.