

Heterogeneous uptake of the C₁ to C₄ organic acids on a swelling clay mineral

C. D. Hatch, R. V. Gough, and M. A. Tolbert

University of Colorado, Department of Chemistry and Biochemistry and the Cooperative Institute for Research in Environmental Sciences, CIRES Room 318, Boulder, Colorado 80309, USA

Received: 10 May 2007 – Published in Atmos. Chem. Phys. Discuss.: 24 May 2007

Revised: 7 August 2007 – Accepted: 19 August 2007 – Published: 24 August 2007

Abstract. Mineral aerosol is of interest due to its physiochemical impacts on the Earth's atmosphere. However, adsorbed organics could influence the chemical and physical properties of atmospheric mineral particles and alter their impact on the biosphere and climate. In this work, the heterogeneous uptake of a series of small organic acids on the swelling clay, Na-montmorillonite, was studied at 212 K as a function of relative humidity (RH), organic acid pressure and clay mass. A high vacuum chamber equipped with a quadrupole mass spectrometer and a transmission Fourier transform infrared spectrometer was used to detect the gas and condensed phases, respectively. Our results show that while the initial uptake efficiency was found to be independent of organic acid pressure, it increased linearly with increasing clay mass. Thus, the small masses studied allow access to the entire surface area of the clay sample with minimal effects due to surface saturation. Additionally, results from this study show that the initial uptake efficiency for butanoic (butyric) acid on the clay increases by an order of magnitude as the RH is raised from 0% to 45% RH at 212 K while the initial uptake efficiency of formic, acetic and propanoic (propionic) acids increases only slightly at higher humidities. However, the initial uptake efficiency decreases significantly in a short amount of time due to surface saturation effects. Thus, although the initial uptake efficiencies are appropriate for initial times, the fact that the uptake efficiency will decrease over time as the surface saturates should be considered in atmospheric models. Surface saturation results in sub-monolayer coverage of organic acid on montmorillonite under dry conditions and relevant organic acid pressures that increases with increasing humidity for all organic acids studied. Additionally, the presence of large organic acids may slightly enhance the water content of the clay above 45% RH. Our results indicate that heterogeneous uptake of or-

ganic acids on swelling clay minerals provides an important irreversible heterogeneous sink for these species.

1 Introduction

Mineral aerosol in the Earth's atmosphere is of interest because of its direct and indirect effect on climate as well as its potential importance for heterogeneous chemistry (Sokolik et al., 1998; Zhang and Carmichael, 1999; Tang et al., 2004). Mineral aerosol components are variable depending on the mineralogy of the source region and atmospheric processing during transport. Recent field measurements have shown that atmospheric mineral aerosol is predominantly composed of layered silicates. For example, Reid et al. (2003) found that 70% of all mineral particle mass originating from Africa consisted of layered silicates such as illite, kaolinite and montmorillonite clays. Montmorillonite is of particular interest because it has a unique ability to accommodate an aqueous environment and provide a large reactive surface area for adsorption. Additionally, this clay is known to contain comparable quantities of water to common deliquescent species, such as ammonium sulfate (Frinak et al., 2005). Thus, the hygroscopic nature of montmorillonite may provide an excellent platform for uptake of gas phase species in the Earth's troposphere and may make it a good cloud nucleus.

As mineral particles are transported through the atmosphere, the particles encounter many environments. Mineral aerosol is generally emitted from dry regions where humidity is low. However, the emitted particles may be entrained in the troposphere for days to weeks (Bauer et al., 2004). For example, an estimated 50% of airborne mineral particles from Chinese deserts are transported thousands of kilometers over the remote North Pacific Ocean (Zhang et al., 1997). Additionally, the entrained dust has been found to reach the United States (DeBell et al., 2004) and Canada (McKendry et al., 2001). As the mineral aerosol moves out of the source

Correspondence to: C. D. Hatch
(courtney-hatch@uiowa.edu)

region it can pass through semi-arid regions where biomass burning often occurs, providing a substantial source of organics to the atmosphere. Additionally, transport over urban regions can result in the mixing of pollution with dust plumes (Maxwell-Meier et al., 2004; Sun et al., 2005; Wang et al., 2005). Mineral particles en route through these regions could have a significant impact on the gas phase abundance of species such as nitric acid, sulfuric acid and organics and may help transport these species over longer distances. Further, swelling clays, such as montmorillonite, can provide a reactive environment for heterogeneous chemistry under more humid conditions as the particles are transported out of the arid source region.

Field measurements have shown that mineral particles entrained in the atmosphere often contain organic components. Single particle analysis of mineral aerosol has shown that oxygenated organic compounds are found in the same particles as mineral species, including aluminosilicate clays (Haderlein and Schwarzenbach, 1993; Lee et al., 2002; Lee et al., 2003; Falkovich et al., 2004). Russell et al. (2002) used soft X-ray spectromicroscopy to map adsorbed components on atmospheric aerosol particles. This study found a strong interaction between mineral aerosol and carbonyl species, possibly aldehydes, ketones and organic acids. Additionally, Falkovich et al. (2004) studied dust samples collected over an urban region in Israel's coastal plain. Scanning electron microscopy coupled with an energy dispersion system (SEM/EDS) revealed that some of the collected particles were agglomerates of aluminosilicates containing heterogeneous organic coatings. Additional gas chromatography/mass spectrometry and ion chromatography measurements indicated that formic and acetic acids were the most common mono-carboxylic acids found in the mineral aerosol samples. Falkovich et al. (2004) also found that carboxylic acids adsorbed more efficiently to Saharan mineral aerosol particles at higher relative humidity (RH) with respect to liquid water, suggesting water-assisted uptake onto entrained mineral particles. Thus, heterogeneous uptake of organic acids appears to play a role in the atmospheric processing of mineral aerosol and may be enhanced under humid conditions. However, very little is known about the kinetics and mechanisms of these interactions.

A few laboratory studies have begun to explore the heterogeneous chemistry of organic compounds with various types of minerals (Li et al., 2001; Al-Abadleh and Grassian, 2003; Carlos-Cuellar et al., 2003; Al-Hosney et al., 2005). Previous studies probing the uptake of acetic acid at 295 K on minerals such as Al_2O_3 , $\alpha\text{-Fe}_2\text{O}_3$ and SiO_2 and formic acid at 295 K on CaCO_3 have been reported (Al-Hosney et al., 2005; Carlos-Cuellar et al., 2003). Al-Hosney et al. (2005) found that addition of a small amount of water vapor enhanced the affinity of CaCO_3 for formic acid. These laboratory studies have shown that some minerals representative of airborne dust may have a sizable capacity to adsorb small organic acids. However, previous studies have not yet

quantified the heterogeneous uptake of organic acids under typical upper tropospheric temperatures and atmospherically relevant RH values, nor have they explored the aqueous environment of a swelling clay which may impact gas-to-particle conversion of organic acids in the atmosphere.

Previous studies in our laboratory have explored the heterogeneous adsorption of water and nitric acid on Na-montmorillonite clay (Frinak et al., 2005; Mashburn et al., 2006). These studies have suggested that swelling clays may be important cloud condensation nuclei because of their large water adsorption capacities. Additionally, in the presence of nitric acid, montmorillonite can accommodate even more water at RH values greater than 30% (Mashburn et al., 2006). Thus, in addition to the potential effects on gas phase concentrations of organic acids, the hygroscopic properties of these clays may also be altered in the presence of organics. These previous studies have allowed for a detailed characterization of Na-montmorillonite as a function of RH. The present study explores the heterogeneous uptake of the C_1 to C_4 organic acids, including formic, acetic, propanoic and butanoic acids, at upper tropospheric temperatures on Na-montmorillonite clay as a function of RH, organic acid pressure and clay mass.

2 Experimental methods

2.1 Vacuum chamber

The Knudsen cell flow chamber used for the present study has been described in detail previously and is briefly discussed here (Mashburn et al., 2006). A high vacuum flow reactor equipped with a UTI 100C quadrupole mass spectrometer and a Nicolet 740 Fourier transform infrared (FT-IR) spectrometer in transmission mode is used to simultaneously monitor the gas and condensed phases, respectively. The stainless steel chamber is evacuated through an escape orifice, 0.16 cm^2 , by a 210 L/sec turbomolecular pump (Pfeiffer) reaching baseline pressures near 2.7×10^{-7} hPa. The gas phase organic acid and water vapor are introduced through two separate leak valves backed by the vapor pressure above the liquid phase held at 273 K in an ice bath. The organic acids are housed in glass bulbs and are degassed by daily freeze-pump-thaw cycles. The design of this vacuum chamber allows for isolation of the reactive surface from the gas phase flow by covering the sample with a retractable Teflon cup that seals against the chamber walls with an O-ring. Thus, a steady-state flow of gas phase reactants may be established prior to exposure of the clay.

An MKS Baratron capacitance manometer and an ionization gauge measure the total chamber pressure while the absolute pressure of the organic acid is monitored by mass spectrometry. Prior to each experiment, the mass spectrometer signal for the organic acid is calibrated to the partial pressure of the acid measured by the Baratron and ionization

gauge. The partial pressure of the acid is obtained from the mass spectrometer signal after equilibration with the water vapor at the experimental temperature and prior to exposure to the clay sample. For the experiments reported here, organic acid pressures ranged from 8×10^{-6} to 1.3×10^{-4} hPa in order to maximize signal while staying in the molecular flow regime. Uncertainty in the organic acid partial pressure is less than 5% for all organic acids studied based on uncertainty in the mass spectrometer signal and the calibration curve collected prior to each experiment. The partial pressure of water, obtained by subtracting the partial pressure of the organic acid from the total chamber pressure, is used to determine the RH at the temperature of the sample (Flatau et al., 1992). Based on the error in the sample temperature and partial pressure of water vapor, the uncertainty in the RH is less than 10% under humid conditions. The water pressure is maintained below the saturation vapor pressure with respect to water ice for all experiments reported here.

The clay sample is deposited onto an infrared-transparent silicon wafer and the wafer is attached to a gold-coated copper mount with stainless steel clips. Indium foil is placed between the sample and mount to improve thermal contact. Upper tropospheric temperatures are reached using a liquid nitrogen-cooled cryostat. The temperature is set by resistively heating against the liquid nitrogen using a Watlow heater controlled by an 818 Eurotherm temperature controller. Two Type T thermocouples attached to either side of the sample monitor the temperature of the sample. The temperature of the thermocouple is calibrated by performing ice frost point calibrations (Marti and Mauersberger, 1993) and is held at 212 K for all experiments reported here.

2.2 Sample preparation

The chemical composition, surface area and infrared analysis of Na-montmorillonite, SWy-2, clay obtained from the Clay Mineral Society's Clay Source Repository, have been reported previously (Van Olphen and Fripiat, 1979; Madejova and Komadel, 2001; Frinak et al., 2005). The clay sample is prepared by depositing a well-mixed slurry of 100 mg SWy-2 in 25 mL water (Fisher Scientific, HPLC grade) on the silicon wafer. The deposited slurry is allowed to dry under ambient conditions and is then dried further by evacuation in the vacuum chamber for 24 to 48 h or until a background pressure of approximately 2.7×10^{-7} hPa is reached. This method results in consistent sample masses near 1.5 mg and geometric surface areas near 2 cm^2 .

Transmission (TEM) and scanning electron microscopy (SEM) were used to examine the SWy-2 particle size and surface characteristics. Figure 1a shows a representative TEM image of the SWy-2 clay, where the particles are observed as the dark regions. This sample was prepared with a very dilute solution of the clay in water and yielded much smaller total masses than those used in the experiments. As shown, the clay particles are approximately $2 \mu\text{m}$ in diameter and




Fig. 1. (a) TEM and (b) SEM images of SWy-2 particles.

are non-spherical. Figure 1b shows an SEM image of SWy-2 prepared in a similar manner as described for the uptake experiments. This image shows the rough surface morphology of the clay particles under experimental conditions.

2.3 Experimental procedure

During an experiment, an FT-IR background scan of the dry clay is first collected at the experimental temperature. The Teflon cup is then closed and the flow of organic acid is established through a leak valve. The chamber walls are passivated until a steady flow is achieved, approximately 45 min. For experiments performed under humid conditions, water vapor is added using a separate leak valve after sufficient passivation of the walls with the organic vapor. The experimental partial pressure of the organic species is obtained from the mass spectrometer after equilibration with the water and organic vapors for approximately 45 additional minutes. Once

a constant flow of the acid and water vapor is established, the initial uptake is measured by opening the cup and exposing the sample to the gas phase for 200 s. Additional experiments at longer exposure times allowed for characterization of the condensed phase by FT-IR spectroscopy as the clay fully expands upon water adsorption. Our measurements were limited to studying the uptake of the organic acids on SWy-2 upon simultaneously exposing the samples to the gas phase water and organic acid because the surface area of the clay under swollen conditions is unclear and the time for the clay to reach equilibrium with the water vapor exceeds six hours. Although we cannot obtain an accurate initial uptake efficiency from experiments in which the clay is sequentially exposed first to gas phase water and then to the organic acid vapor because we do not know the clay surface area under the swollen conditions, as discussed below, FT-IR measurements do indicate that formic and acetic acid were taken up by the previously swollen wet clay.

2.4 FT-IR data analysis

Infrared spectra of the clay upon exposure to the gas phase were obtained when the Teflon cup was open by averaging 32 scans at 4 cm^{-1} resolution and were ratioed to the background spectrum of the clay collected prior to each experiment. For each RH, the adsorbed products were monitored by transmission FT-IR spectroscopy as a function of time until the condensed phase was no longer observed to change, typically for three to five hours. Blank experiments, in which no clay sample was present, were performed for the same conditions and exposure times as experiments where the clay was present on the silicon wafer. No changes were observed in the infrared spectra for blank experiments under all conditions studied. FT-IR measurements allowed for quantification of adsorbed water as described in previous papers (Frinak et al., 2005; Mashburn et al., 2006).

2.5 MS data analysis

2.5.1 Uptake efficiency measurements

Mass spectrometer measurements with two second time resolution allowed for direct observation of changes in the gas phase organic acid concentration during each experiment. The initial drop in the mass spectrometer signal of the organic acid upon opening the cup reaches a minimum in the first two seconds of exposure, followed by recovery due to surface saturation. Thus, the mass spectrometer signal upon opening the Teflon cup used for the calculation of the initial heterogeneous uptake efficiency, γ_o , was collected at two seconds after exposure. The uptake efficiency is defined as the ratio of the number of gas phase molecules lost to the surface per unit time to the total number of collisions with the surface per unit time. The γ_o is obtained from mass spectrometer data in the same manner as was determined pre-

viously for nitric acid uptake on SWy-2 and shown below (Mashburn et al., 2006):

$$\gamma_o = \frac{A_h}{(\text{SSA}_{\text{BET}} \times m)} \left(\frac{I_o - I_t}{I_t} \right) \quad (1)$$

In Eq. (1), SSA_{BET} is the specific surface area (cm^2/mg) of SWy-2, A_h is the effective area of the escape orifice, $A_h=0.16\text{ cm}^2$, calibrated using nitrogen gas and m is the mass of the SWy-2 sample. I_o is the initial mass spectrometer signal of the organic acid before opening the Teflon cup to expose the sample and I_t denotes the mass spectrometer signal of organic acid after two seconds of exposure, corrected for blank experiments performed under the same conditions. For all experiments reported here, initial uptake on the blank silicon wafer is significantly less than uptake observed when the SWy-2 sample is present. A correction for blank experiments introduces an uncertainty of approximately 7% into the calculation of γ_o , based on the reproducibility of blank measurements. Additionally, independent mass spectrometer measurements scanned the gas phase for potential products during exposure of the organic acids to SWy-2. Under all conditions studied, the evolution of gas phase products during exposure was not observed.

2.5.2 Clay surface area

The swelling properties of montmorillonite clay have been studied for many decades and a more detailed description of water adsorption and swelling may be found in the literature (Mooney et al., 1952; Anderson et al., 1978; Hall and Astill, 1989; Cases et al., 1992; Frinak et al., 2005). In the absence of water vapor, only the external surfaces of montmorillonite are available for adsorption. Water begins to penetrate the interlayer space of the clay above 16% RH and reaches monolayer coverage of water, both internal and external, near 48% RH (Newman, 1983; Cases et al., 1992; Hensen and Smit, 2002). Water begins adsorbing in multiple layers at higher humidities. Although the swelling regimes of montmorillonite clay have been experimentally investigated at room temperature as a function of RH by Cases et al. (1992) and Newman (1983), previous studies in our laboratory (Frinak et al., 2005) have shown that water adsorption on montmorillonite clay is independent of temperature and thus these regimes also apply at upper tropospheric temperatures. Upon swelling, access to both internal and external surfaces results in the availability of a significantly larger surface area compared to the dry external surface area (Brunauer et al., 1938; Cases et al., 1992; Frinak et al., 2005).

To determine γ_o , the appropriate surface area is needed. As shown in the SEM image of the SWy-2 particles, the rough morphology of the clay particles implies that the geometric surface area, assuming a flat surface, is a significant underestimate of the available surface area. Non-polar molecules, such as nitrogen, are known to only adsorb to the

external surfaces of montmorillonite and thus provide a direct measurement of the external surface area including clay microstructure (Hendricks et al., 1940; Mooney et al., 1952; Pennell et al., 1992). The BET surface area for SWy-2 is $318.2(\pm 2.2)$ cm²/mg (Dogan et al., 2006), as reported by the Clay Mineral Society and independently confirmed by the authors by BET analysis using a volumetric expansion technique with nitrogen as the adsorbate. The calculated BET surface area was within the uncertainty reported by the Clay Mineral Society. However, when swollen, the available surface area of SWy-2 can increase as much as 25 times that of the external surface area obtained from BET measurements (Frinak et al., 2005; Mashburn et al., 2006).

Previous studies have determined the surface area of the swollen clay at various RH values using gravimetric (Cases et al., 1992) and FT-IR techniques (Frinak et al., 2005) to probe water adsorption. Frinak et al. (2005) found that saturation of adsorbed water on SWy-2 at sufficiently low water pressures required long exposure times. For example, reducing the water pressure from 1.3×10^{-2} to 4×10^{-3} hPa increases the $1/e$ loading time by a factor of four, from approximately 10 min to 40 min. The measurements reported here were conducted under low water pressure conditions; therefore, we expect very long exposure times to be required for the SWy-2 clay to become swollen and the internal surfaces accessible. Thus, the interlayer space of the clay is not accessible to the organic acids at the early time of two seconds and only the external surfaces are available for adsorption during initial uptake (Frinak et al., 2005; Mashburn et al., 2006). As discussed in further detail later, surface saturation and swelling effects have a minimal impact on the γ_o measurement. Additionally, the availability of the entire BET surface area is confirmed by studying the mass dependence of the heterogeneous uptake of formic, acetic, propanoic and butanoic acids on SWy-2 at 212 K and 29% RH.

2.6 Quantification of adsorbed organic acid

Mass spectrometer measurements may also be used to quantify the total amount of organic acid adsorbed to SWy-2 (Frinak et al., 2004). The organic acid content adsorbed to SWy-2 is determined by integrating the area under the mass spectrometer signal while the sample is exposed after correction for loss to the sample mount by subtraction. The mass spectrometer signal was first converted to a flow (molecules/sec) using the Knudsen effusion relation:

$$F = \frac{P_{\text{org}} A_h}{(2\pi M_w k T)^{1/2}} \quad (2)$$

where A_h is the effective area of the escape orifice, 0.16 cm², to the turbomolecular pump, M_w is the molecular weight of the organic acid, k is the Boltzmann constant and T is the temperature of the gas phase, defined as room temperature. The flow of organic acid during exposure to the clay is then offset and inverted so that the area under a double

Table 1. Identification of FT-IR peaks observed during exposure of organics to SWy-2 clay.

Peak Position (cm ⁻¹)	Assignment
~1620	Dissociated carbonyl species ^{a,b}
1350 – 1475	Deformations of CH ₃ groups ^a
3624	Structural OH stretch ^c
3350	OH stretch ^c
1640	HOH bend ^c

^a from Grassian (2001)

^b from Max and Chapados (2004) and Kubicki et al. (1999)

^c from Bishop et al. (1994)

exponential fit through the uptake data gives the number of organic acid molecules adsorbed to the surface (Al-Abadleh and Grassian, 2000; Frinak et al., 2004). Similar to previous studies (Al-Abadleh and Grassian, 2000; Frinak et al., 2004), we have found that a double exponential fit represents the shape of the uptake curve more accurately than a single exponential fit. The number of organic acid molecules adsorbed at steady-state, n_{org} , is obtained by extrapolating the data collected during a 200 s experiment to 5000 s. Extrapolation to 5000 s results in less than 10% increase in the calculated organic acid content values. However, extrapolation beyond 5000 s does not change the calculated steady-state organic acid content. Thus, this extrapolation procedure is sufficient to obtain realistic coverage values on the montmorillonite clay. The n_{org} is then corrected for organic acid uptake on the silicon wafer, mount and chamber walls when the clay sample was not present. Under all conditions studied, less than 20% of the organic acid lost from the gas phase is taken up by the sample wafer, mount and chamber walls. Finally, n_{org} can then be converted to a mass and divided by the mass of the clay sample resulting in a mass ratio giving the organic acid content in units of mg_{org}/g_{clay}. Uncertainty in the organic acid content due to blank experiments is approximately 9%, based on the reproducibility of blank measurements. Alternatively, the number of adsorbed organic acid molecules can be divided by the surface area of the SWy-2 clay to estimate the acid coverage (molec/cm²). Isothermal desorption is also measured to determine the reversibility of the interaction.

3 Results

3.1 FT-IR measurements of condensed products

FT-IR measurements revealed that under dry conditions, even after five hours of exposure to organic acids, no adsorbed species were observed in the infrared spectra. This suggests that the organic acids do not have a strong affinity for SWy-2 when water vapor is not present and the clay is not




Fig. 2. FT-IR spectra collected (a) before and (b) after exposure of a 1.2 mg SWy-2 sample to 1.3×10^{-4} hPa acetic acid at 212 K and 29% RH. (c) Also shown is the infrared spectrum collected after warming the sample to room temperature to drive off adsorbed water. All spectra are ratioed to a background spectrum of the clay collected prior to exposure at the same temperature.

swollen. However, in the presence of water vapor, there is evidence for organic uptake. Figure 2 shows the FT-IR spectra collected before and after exposure of acetic acid to SWy-2 at 212 K and 29% RH. FT-IR peak assignments based on previous studies are shown in Table 1 (Bishop et al., 1994; Grassian, 2001; Shilling and Tolbert, 2004; Shilling et al., 2006). In the presence of acetic acid and water vapor, the most prominent features observed after exposure are the stretch and bend of adsorbed water at 3350 cm^{-1} and 1640 cm^{-1} , respectively (Bishop et al., 1994). Additionally, a small change in the structural OH stretching region at 3624 cm^{-1} is observed. This region represents stretching of the hydroxyl groups bound directly to the aluminosilicate structure of the montmorillonite clay. All of these features are typical of an experiment in which SWy-2 is exposed only to water without any organic acid present (Frinak et al., 2005). However, since the dominant feature of carbonyl species, the carbonyl stretch, is normally found near 1700 cm^{-1} , an overlap with the water bending peak would occur.

Previous studies have shown that adsorbed water is completely reversible upon warming the SWy-2 sample to room temperature (Frinak et al., 2005). Thus, following each experiment, adsorbed water was removed by warming to room temperature and the resulting infrared spectrum is shown in Fig. 2. Upon removal of adsorbed water, a broad feature, from 1350 to 1470 cm^{-1} , and a peak near 1615 cm^{-1} remain. These peaks are observed for all organic acids studied at experiments performed at 29% and 45% RH after warming to room temperature. The broad feature near 1400 cm^{-1} is attributed to the methyl deformation of adsorbed organic species (Grassian, 2001). The peak near 1615 cm^{-1} is not apparent in the infrared spectrum collected when water is

present, probably due to overlap with the water bend. Additionally, this peak was not observed in previously reported water only measurements (Frinak et al., 2005) nor was it observed in the infrared spectrum of dry SWy-2 reported by the Clay Mineral Society (Madejova and Komadel, 2001). Previous studies showing adsorption of the C₁ to C₄ organic acids to ammonium nitrate found that the organic acid ionizes on the surface and the carbonyl peak, normally observed near 1700 cm^{-1} , shifts to lower wavenumbers near 1580 cm^{-1} based on literature spectra of sodium acetate (Kubicki et al., 1999; Max and Chapados, 2004; Shilling and Tolbert, 2004; Shilling et al., 2006). This result is also in agreement with bulk studies of acetic acid on montmorillonite clay (Kubicki et al., 1999). Thus, the infrared peak near 1615 cm^{-1} is probably due to carboxylate species on the surface or possibly trapped in the interlayer spaces of the clay, such as formate and acetate. We also find that after heating and evaporation of water, the infrared absorption band of the remaining carboxylate peak is similar in strength whether water and the organic acid are adsorbed simultaneously or sequentially. However, it is unclear as to whether dissociation occurs under humid conditions when adsorbed water is present and the clay is swollen, or if the carboxylic acid dissociates upon removal of adsorbed water when warming the sample to room temperature. Regardless of the mechanism for dissociation, results from infrared analysis suggest irreversible uptake of organic acids on SWy-2 resulting in a change in the chemical composition of the clay. Thus, atmospheric processing through heterogeneous uptake of short-chain mono-carboxylic acids and transport through varying humidity conditions can result in modified chemical composition of the particles which may impact the optical properties and further reactivity toward other trace gases in the atmosphere.

Infrared measurements also allowed us to probe the effect of organic acids on water adsorption by the clay. The water content, in units of $\text{g}_{\text{H}_2\text{O}}/\text{g}_{\text{clay}}$, at steady-state is determined by FT-IR spectroscopy, as in previous studies (Frinak et al., 2005; Mashburn et al., 2006), using the optical constants for bulk water (Downing and Williams, 1975). Figure 3 shows the steady-state water content of SWy-2 at 212 K in the presence of the organic acids at 29% and 45% RH compared to water only measurements performed at 222 K as a function of RH (Frinak et al., 2005). The error bars represent the variability in the data based on multiple experiments performed under similar conditions. Only the OH stretching region observed near 3350 cm^{-1} was used for this calculation due to possible interference of the carbonyl stretch with the water bend at 1640 cm^{-1} . Although the water only measurements were performed at 222 K, water adsorption is expected to be independent of temperature (Frinak et al., 2005). Thus, the 10 K difference in temperature between the water content in the presence of organic acids and that when the organic acids are not present should have no effect on the water content values reported in Fig. 3. Figure 3 shows that, at 29% RH, the water content does not appear to deviate from the value




Fig. 3. Steady-state water content of SWy-2 as a function of relative humidity. The open circles represent water only experiments performed at 222 K (Frinak et al., 2005) and the closed symbols represent the water content adsorbed on SWy-2 at 212 K in the presence of the organic acids. The inset shows the enhanced water content for propanoic and butanoic acids on an expanded scale.

obtained when no organic is present. A previous study of water adsorption in the presence of nitric acid observed similar results (Mashburn et al., 2006). However, this study also found that nitric acid only influenced the water content above 30% RH. Similar to the nitric acid study, Fig. 3 shows that, at 45% RH, an enhancement in the adsorbed water content is observed in the presence of propanoic (triangle) and butanoic (bowtie) acids by almost a factor of two compared to water only measurements (Frinak et al., 2005). The inset of Fig. 3 shows the enhanced water content at 45% RH on an expanded scale. As measurements of the water content above 45% RH were not attainable in the experimental system, we cannot be certain that this trend will continue at higher humidities. However, nitric acid adsorption showed continuously higher water contents above 30% RH compared to water only measurements (Mashburn et al., 2006). The results reported here may indicate that the same trend will occur as the RH is raised above 45% RH and we could expect more adsorbed water at higher RH values compared to water only measurements. However, more studies are needed at higher RH values to validate this assumption. Additionally, the organic acid partial pressures used in our experiments are much larger than those that prevail in the atmosphere; therefore, organic acids may have less of an impact on water adsorption in the Earth's atmosphere.

3.2 MS measurements

3.2.1 Isothermal desorption to test for reversibility

Isothermal desorption measurements were performed to probe the reversibility of the heterogeneous interaction of organic acids with SWy-2. Upon completion of each uptake




Fig. 4. Raw uptake curves of propanoic acid on 1.6, 5.7, 7.8 and 17.1 mg SWy-2 samples at 212 K, 29% RH and approximately 1.3×10^{-4} hPa propanoic acid. The inset shows an expanded view of the propanoic acid uptake curve on the 1.6 mg SWy-2 clay sample. The vertical lines represent the time SWy-2 was exposed to the gas phase. The traces are offset for clarity.

experiment, while the cup was still closed, the flow of organic acid was discontinued and the vacuum chamber was evacuated to the baseline pressure. Once a stable baseline pressure was established, the Teflon cup was retracted and isothermal desorption of the organic acid from SWy-2 was monitored by mass spectrometry. Integration of the area under the desorption peak resulted in the number of desorbed molecules. Upon comparing to the number of molecules adsorbed during the experiment we found that less than 20% of the adsorbed organic acid desorbed upon opening the cup for all of the organic acids and under all conditions studied. However, as mentioned previously, up to 20% of the organic acid lost from the gas phase was due to adsorption on the sample wafer, mount and chamber walls. Thus, it is possible that some of the desorbed organic acid remains on the exposed wafer and mount and would therefore not be measured by the mass spectrometer. This would result in an overestimate of the irreversibility of organic acid uptake. However, the fact that an organic spectral signature remains on the clay even after warming to room temperature indicates that the adsorption is at least partially irreversible.

3.2.2 Uptake as a function of clay mass

Mass spectrometry was used to directly monitor loss of the organic acid to the clay surface as a function of clay mass. Figure 4 shows the raw uptake curves obtained from the mass spectrometer during exposure of 1.6, 5.7, 7.8 and 17.1 mg SWy-2 samples to approximately 1.3×10^{-4} hPa propanoic acid at 212 K and 29% RH. The inset of Fig. 4 shows an expanded view of the propanoic acid uptake curve on a 1.6 mg SWy-2 sample. Similar traces are observed for the other organic acids studied. These uptake curves show that the




Fig. 5. $(I_o - I_t)/I_t$ obtained from organic acid uptake measurements at 212 K and 29% RH as a function of SWy-2 mass after two seconds of exposure. Formic, acetic and propanoic acid uptake experiments were performed in the presence of approximately 1.3×10^{-4} hPa acid while butanoic acid uptake measurements were performed in the presence of approximately 1.3×10^{-5} hPa butanoic acid.

uptake of organic acids on small SWy-2 samples appears to reach saturation within the 200 s time scale of exposure, indicated by complete recovery, within error, of the mass spectrometer signal to baseline values. The incomplete recovery of the mass spectrometer for the larger sample masses prior to closing the cup may indicate diffusion into and adsorption on the underlying clay layers at longer exposure times. Since larger sample masses show incomplete recovery of the mass spectrometer signals, longer exposure times are required to accurately determine the organic acid coverage at steady-state. Thus, the organic acid content is more accurately known for the small sample masses used in the RH and pressure studies, discussed below.

To determine the appropriate surface area for quantification of γ_o , the heterogeneous uptake of the C₁ to C₄ organic acids was studied as a function of SWy-2 mass at 212 K, 29% RH and approximately 1.3×10^{-4} hPa organic acid, 1.3×10^{-5} hPa for butanoic acid. Figure 5 shows $(I_o - I_t)/I_t$, where I_t is measured after two seconds of exposure and corrected for blank experiments, as a function of SWy-2 mass for each of the organic acids studied. As expected from previous reports, $(I_o - I_t)/I_t$ values appear to increase linearly for small sample masses (Carlos-Cuellar et al., 2003; Al-Hosney et al., 2005; Mashburn et al., 2006). Sample masses less than




Fig. 6. Initial uptake efficiencies for (a) formic, (b) acetic, (c) propanoic and (d) butanoic acid uptake on approximately 1.5 mg SWy-2 at 212 K and 0% RH as a function of organic acid pressure. The horizontal lines represent the average γ_o value obtained over the range of partial pressures studied.

5 mg were considered to lie within the linear mass dependent regime. For masses larger than 5 mg, $(I_o - I_t)/I_t$ appears to saturate, suggesting that all of the underlying clay layers are not accessible by the adsorbate. Thus, the entire sample mass is available for adsorption on the time scale of the measurement if the samples remain smaller than 5 mg, such as those used in the RH and pressure studies (~ 1.5 mg). Thus, the BET surface area is used to determine the γ_o from these measurements. The slope obtained from a linear fit through the linear mass dependent region is used to calculate an average initial uptake efficiency, γ_o^{avg} , over the mass range studied as follows:

$$\gamma_o^{\text{avg}} = (\text{slope}) \frac{A_h}{SSA_{\text{BET}}} \quad (3)$$

As shown in Table 2, we find γ_o^{avg} values of $1.7(\pm 0.3) \times 10^{-5}$, $2.0(\pm 0.4) \times 10^{-5}$, $3.2(\pm 0.2) \times 10^{-5}$ and $4.0(\pm 0.1) \times 10^{-5}$ for the heterogeneous uptake of formic, acetic, propanoic and butanoic acids, respectively, on SWy-2 at 212 K and 29% RH from the data shown in Fig. 5.

3.2.3 Uptake as a function of organic acid pressure


Results from the mass study are based on organic acid pressures near 1.3×10^{-4} hPa to maximize signal. This value is much higher than the partial pressure of organic acids found

Table 2. The initial uptake efficiencies for uptake of formic, acetic, propanoic and butanoic acids on SWy-2 at 212 K as a function of RH from 0% to 45% RH. The limit of detection for these measurements is $\gamma_o^{\text{LOD}}=1 \times 10^{-6}$.

Organic	0% RH	29% RH	45% RH
Formic acid	1.7×10^{-5}	1.7×10^{-5}	2.3×10^{-5}
Acetic acid	1.3×10^{-5}	2.0×10^{-5}	6.0×10^{-5}
Propanoic acid	5.4×10^{-5}	3.2×10^{-5}	6.2×10^{-5}
Butanoic acid	2.2×10^{-5}	4.0×10^{-5}	2.2×10^{-4}

in the troposphere. Previous studies have shown that γ_o can vary significantly with pressure (Underwood et al., 2000). We have measured the γ_o values for the C₁ to C₄ organic acids on SWy-2 as a function of organic acid pressure from 8×10^{-6} to 1.3×10^{-4} hPa at 212 K and 0% RH. The initial uptake efficiency was determined using Eq. (1) assuming the external BET surface area of the clay is available for uptake. Figure 6 shows γ_o obtained from mass spectrometer data for organic acid uptake on approximately 1.5 mg SWy-2 at 212 K and 0% RH as a function of organic acid pressure. Error in the initial uptake efficiency represents the average uncertainty, $\pm 36\%$, due to measurement reproducibility of uptake on the clay and blank wafer under similar conditions. The horizontal lines represent the average γ_o obtained over the range of pressures studied. Similar to a previous study of acetic acid on $\alpha\text{-Fe}_2\text{O}_3$ (Carlos-Cuellar et al., 2003), we find γ_o to be independent of organic acid pressure for all of the acids studied. Therefore, gas phase adsorption of organic acids is a first-order process within the range of pressures studied. The average γ_o for heterogeneous uptake of formic, acetic, propanoic and butanoic acids on approximately 1.5 mg SWy-2 at 212 K and 0% RH are $1.7(\pm 0.6) \times 10^{-5}$, $1.3(\pm 0.5) \times 10^{-5}$, $5.4(\pm 1.9) \times 10^{-5}$ and $2.2(\pm 0.8) \times 10^{-5}$, respectively. These values are reported in Table 2.

Organic acid coverage on SWy-2 is determined from the mass spectrometer data as described previously and corrected for blank experiments performed under the same conditions. Figure 7 shows the steady-state coverage of formic, acetic, propanoic and butanoic acids on 1.5 mg SWy-2 at 212 K and 0% RH as a function of organic acid pressure. We find that the amount of adsorbed organic acid increases with acid pressure. Sub-monolayer coverages of formic and acetic acids are found to increase linearly with organic acid pressure, never reaching saturation within the range of pressures studied. However, butanoic and perhaps propanoic acids appear to saturate at approximately one monolayer for pressures greater than 2.7×10^{-5} hPa. Thus, under dry conditions, sub-monolayer coverage of organic acid is expected at relevant vapor pressures on atmospheric mineral aerosol surfaces.

**Fig. 7.** The steady state coverage of (a) formic, (b) acetic, (c) propanoic and (d) butanoic acids on 1.5 mg SWy-2 at 212 K and 0% RH as a function of organic acid pressure.

3.2.4 Uptake as a function of relative humidity

Previous studies have shown that the C₁ to C₆ carboxylic acids favor the particle phase under humid conditions based on the Henry's law constant for these acids in water (Khan et al., 1995). Thus, under humid conditions, it is probable that swelling clay minerals may be a significant sink for organic acids. The heterogeneous uptake efficiency of the C₁ to C₄ organic acids on 1.5 mg SWy-2 samples was measured as a function of RH from 0% to 45%, at 212 K and approximately 1.3×10^{-4} hPa organic acid, 1.3×10^{-5} hPa for butanoic acid. The measured γ_o values are summarized in Table 2 and shown graphically in Fig. 8a. At 0% RH, γ_o is independent of pressure and thus the average values obtained from the pressure study are reported here. The γ_o values at 29% RH represent γ_o^{avg} determined from the mass study. Assuming that the organic acids have access to the




Fig. 8. (a) The initial uptake efficiencies on a log scale for the uptake of formic, acetic, propanoic and butanoic acids on approximately 1.5 mg SWy-2 at 1.3×10^{-4} hPa organic acid pressure, 1.3×10^{-5} hPa butanoic acid and 212 K as a function of RH. (b) The organic acid content, $\text{mg}_{\text{org}}/\text{g}_{\text{clay}}$, of adsorbed formic, acetic, propanoic and butanoic acids on approximately 1.5 mg SWy-2 at 212 K as a function of RH. Lines are drawn to guide the eye.

entire sample mass over the range of humidities studied, γ_o at 45% RH was also determined. Error in the reported γ_o values shown in Fig. 8a are based on the combined uncertainty from the reproducibility of uptake on the clay and blank silicon wafer. Figure 8a shows that γ_o does not appear to vary significantly at RH values less than 29%. However, we find that the initial uptake efficiency of butanoic acid increases by nearly an order of magnitude as the RH was raised from 0% to 45%. This may be due to enhanced acid dissociation on the clay surface under humid conditions, as suggested by infrared analysis. However, the uptake of formic, acetic and propanoic acids increase only slightly with increasing RH. Thus, adsorbed water appears to influence the initial uptake efficiency of some organic acids more so than others, particularly near 45% RH. Higher humidities may result in even larger initial uptake efficiencies. This represents the first report of the γ_o for small organic acids on smectite clay under tropospherically relevant humidities and temperatures.

Organic acid uptake was found to vary with the type of organic acid studied. Under all conditions, formic acid was found to have the lowest reactivity with the SWy-2 surface. To help understand these results, Table 3 shows the number of carbon atoms, molecular weight, boiling point and enthalpy of vaporization for each acid studied (Lide, 2004).

Additionally, the γ_o values at 45% RH are also shown in Table 3. The γ_o appears to follow general trends in molecular weight, boiling point, vapor pressure and the number of carbon atoms in the hydrocarbon chain; all of which influence the initial interaction between the clay surface and gas phase organic acid.

Previous studies of nitric acid uptake on SWy-2 clay as a function of RH found that the γ_o was not measurable below 16% RH (Mashburn et al., 2006). However, above 16% RH, the γ_o for nitric acid uptake on SWy-2 at 221 K increased dramatically with RH. In contrast to nitric acid uptake, measurable uptake was reported here for all of the organic acids at 0% RH. We find that the behavior of butanoic acid with increasing RH is very similar to that observed for nitric acid uptake on the same clay. However, under humid conditions, the γ_o from organic acid uptake is smaller by a factor of two to an order of magnitude from the values observed for nitric acid (Mashburn et al., 2006). Thus, organic acid uptake on smectite clays is less efficient than nitric acid uptake under humid conditions.

Under humid conditions, it is possible for water adsorption to influence the measured initial uptake efficiencies within the times scale of the measurement. Water adsorption on montmorillonite clay is very slow under the experimental conditions and thus the measurement of the initial uptake efficiency from simultaneous exposure of gas phase water and organic acid should not be significantly affected. To explore this further, the FT-IR measurements of water adsorption, based on a time resolution of 11 seconds, on SWy-2 in the presence of 1.3×10^{-4} hPa acetic acid at 29% RH can be used according to the method used by Frinak et al. (2005) and Mashburn et al. (2006) to estimate the amount of water adsorbed to SWy-2 after 11 s of exposure. Using the water bend at 1640 cm^{-1} and assuming one monolayer is $1 \times 10^{15} \text{ molec}/\text{cm}^2$, we estimate that there is less than 8% of a monolayer of water after 11 s of exposure using the external BET surface area. Additionally, using the mass spectrometer data for acetic acid uptake under the same conditions, we estimate that less than 1% of a monolayer of acetic acid would be present after two seconds of exposure. Thus, although more than $\sim 91\%$ of the surface sites are still available for uptake and surface saturation effects will be minimal, occupation of surface sites at the time of initial uptake will result in slightly low initial uptake efficiencies. However, this effect will be even less under dry conditions if less than 1% of a monolayer of the organic acid is adsorbed on the clay surfaces within the first two seconds of exposure of the SWy-2 to the organic acid. Additional error in the initial uptake efficiencies reported here may arise from a possible pressure gradient surrounding the clay surface due to uptake prior to the initial measurement by the mass spectrometer. However, because we are in the molecular flow regime, any pressure gradients in our system should be minimal. In any case, experimental uncertainty may result in an underestimate of the true initial uptake efficiency.

Table 3. Properties of the C₁ to C₄ carboxylic acids, including number of carbon atoms, molecular weight, boiling point and enthalpy of vaporization compared to the heterogeneous initial uptake efficiency for the organic acids on SWy-2 at 212 K.

Organic Acid	# of C atoms	M _w (g/mol)	BP* (K)	ΔH _{vap} * (kJ/mol)	γ _o at 45% RH	γ _o ^{sphere} at 0% RH	γ _o ^{sphere} at 45% RH
Formic	1	46	374	20	2.3×10 ⁻⁵	4.0×10 ⁻⁴	5.4×10 ⁻⁴
Acetic	2	60	391	23	6.0×10 ⁻⁵	3.0×10 ⁻⁴	1.4×10 ⁻³
Propanoic	3	74	414	32	6.2×10 ⁻⁵	1.3×10 ⁻³	1.5×10 ⁻³
Butanoic	4	88	437	41	2.2×10 ⁻⁴	5.1×10 ⁻³	5.1×10 ⁻³

* Handbook of Chemistry and Physics, 73rd Ed. (Lide, 2004)

The organic acid content at steady-state was also determined as a function of RH at 212 K and approximately 1.3×10⁻⁴ hPa organic acid, 1.3×10⁻⁵ hPa for butanoic acid. Figure 8b shows the organic acid content, mg_{org}/g_{clay}, of formic, acetic, propanoic and butanoic acids on approximately 1.5 mg SWy-2 samples at 212 K as a function of RH. The error in the organic acid content, ±32%, represents the combined uncertainty due to the reproducibility of uptake measurements on the clay and blank silicon wafer. It can be seen that, as the RH is increased, the organic acid content increases for all organic acids studied. Thus, while the uptake kinetics are not significantly increased, the equilibrium organic acid content does increase with increasing water vapor. This result is in agreement with water-assisted uptake of organic acids on mineral particles suggested by Falkovich et al. (2004).

The propanoic acid shows the largest increase in adsorbed mass with RH, increasing by a factor of five from 0.3% to almost 2% propanoic acid by mass from 0% to 45% RH, respectively. It is interesting to note that propanoic acid has the highest Henry's law constant of the four acids studied (Khan et al., 1995). This may suggest that more soluble organic acids will adsorb to the swelling clay to a greater extent than less soluble species under humid conditions. Additionally, the trend in coverage matches the trend in pK_a with pK_a going from 3.75 (formic), 4.76 (acetic), 4.83 (butanoic) to 4.86 (propanoic) (Lide, 2004). Thus, the swelling properties of montmorillonite clay provide an aqueous environment under humid conditions that appears to impact the adsorption of oxygenated organics.

4 Discussion

Although no direct comparison of the data reported here can be made to previous studies due to different adsorbates, some statements can be made comparing uptake on other minerals found in the troposphere. Similar to findings from previous studies of formic and acetic acids on carbonate and other mineral oxides (Carlos-Cuellar et al., 2003; Al-Hosney et al.,

2005), our results suggest that the organic acids dissociate on the montmorillonite clay under humid conditions. However, differences are apparent in the measured uptake efficiencies. One previous study investigated the heterogeneous uptake of acetic acid on a series of minerals under dry conditions at room temperature (Carlos-Cuellar et al., 2003). That study found irreversible uptake of acetic acid and reported γ_o values of 2×10⁻³, 2×10⁻³ and 2×10⁻⁴ for α-Fe₂O₃, α-Al₂O₃ and SiO₂, respectively, at 8×10⁻⁶ hPa. These values are one to two orders of magnitude larger than the γ_o for acetic acid on SWy-2, γ_o=1×10⁻⁵, reported here under dry conditions. As shown in Fig. 6, the pressure-independent uptake measurements indicate that this difference is not due to the different pressures used in each study. Additional measurements of acetic acid uptake on SWy-2 at room temperature and dry conditions give a γ_o of 3×10⁻⁶; a factor of four lower than the value obtained at 212 K. Thus, temperature dependence is not responsible for the difference in measured uptake efficiencies. We conclude that the discrepancy in measured initial uptake efficiencies is a result of the different reactivity of the substrates studied. The mineralogy of the reactive surface appears to influence the initial heterogeneous uptake efficiency of organic acids on mineral aerosol in the troposphere, and thus should be considered in atmospheric models.

Previous studies have also reported the steady-state surface coverage for acetic acid on α-Al₂O₃ and SiO₂ under dry conditions to be approximately 8×10¹³ and 7×10¹¹ molec/cm², respectively (Carlos-Cuellar et al., 2003). Based on our results shown in Fig. 7b, we estimate the steady-state coverage of acetic acid on SWy-2 at 8×10⁻⁴ hPa to be near 4×10¹² molec/cm²; on the same order as acetic acid coverage observed on other minerals at room temperature and dry conditions (Carlos-Cuellar et al., 2003). Thus, in contrast to the γ_o, the steady-state coverage of organic acids on mineral aerosol does not appear to depend on the mineralogy of the dust. However, since montmorillonite clay provides an aqueous environment under humid conditions, the organic acid coverage will increase as the RH is raised. More studies of organic acid adsorption on other metal oxides and

non-swelling clays are needed to determine the role mineralogy plays in the adsorbed organic acid content under humid conditions.

5 Atmospheric implications

The heterogeneous removal of organic acids from the atmosphere by uptake onto mineral aerosol must occur on fast enough time scales compared to other removal processes in order to be important in the Earth's atmosphere (Ravisankara, 1997). The dominant removal process of many carboxylic acids is through wet and dry deposition. Although our reported initial uptake efficiencies were measured under conditions where the clay samples were not equilibrated with the reported RH, we use the initial uptake efficiencies for formic, acetic, propanoic and butanoic acids on SWy-2 at 212 K and a mineral aerosol loading of $150 \mu\text{m}^2/\text{cm}^3$ (de Reus et al., 2000), typical of a dust storm, to estimate the lifetime, τ (s), for removal by mineral aerosol in the kinetic limit, assuming the dust plume is entirely composed of montmorillonite, as follows:

$$\tau = \frac{4}{\gamma_o \langle v \rangle SA} \quad (4)$$

where $\langle v \rangle$ (cm/s) is the average velocity of the organic molecule and SA (cm^2/cm^3) is the surface area density of the dust event. We estimate the atmospheric lifetime of these organic acids resulting from heterogeneous removal on swelling clay minerals at 29% RH to be longer than one month at upper tropospheric temperatures. At 45% RH, the atmospheric lifetimes for removal of formic, acetic and propanoic acid are reduced to two weeks, while butanoic acid may be removed within six days. However, the true lifetime for removal is likely to be shorter because the estimated mineral aerosol surface area is based on the diameter of smooth atmospheric mineral spheres.

To account for the differences between the surface area we used to obtain the initial uptake efficiency and the surface area reported in field studies, we convert our uptake efficiencies to a value based on the surface area of spherical particles, SA_{sphere} . Assuming spherical clay particles with a particle diameter of $2 \mu\text{m}$, the SA_{sphere} is estimated using the clay mass and a density of $2.2 \text{ g}/\text{cm}^3$. As shown in Table 3, using SA_{sphere} , we estimate the initial uptake efficiencies, γ_o^{sphere} , for organic acid uptake on SWy-2 to be 4.0×10^{-4} , 3.0×10^{-4} , 1.3×10^{-3} and 5.1×10^{-4} for formic, acetic, propanoic and butanoic acids, respectively, at 0% RH and 212 K. Estimated values of γ_o^{sphere} at 45% RH and 212 K are also shown in Table 3. These values are probably more relevant for use in atmospheric models where spherical particles are often assumed. Using the same method as described above, the lifetime of the organic acids due to heterogeneous removal by mineral aerosol can be obtained using the calculated γ_o^{sphere} values. We find that atmospheric removal of all

the organic acids studied is reduced to less than four days at 0% RH and loss is faster as the RH is raised. However, the initial uptake efficiency is not appropriate for use at longer exposure times due to saturation of surface sites, as demonstrated by the rapid recovery of the mass spectrometer signal in Fig. 4, within approximately one minute. Thus, the lifetime for removal of these organic acids may be much longer than predicted.

To explore the effect of surface saturation on atmospheric dust particles, we can assume that there are two parts per billion by volume (ppbv) acetic acid (Talbot, 1995) in a dust storm of $150 \mu\text{m}^2/\text{cm}^3$ dust (de Reus et al., 2000) at 5 km. Using our measured acetic acid content of $5 \text{ mg}_{\text{org}}/\text{g}_{\text{clay}}$ at 29% RH from Fig. 8b and assuming $2 \mu\text{m}$ diameter particles with a density of $2.2 \text{ g}/\text{cm}^3$, we estimate that approximately 40% of the gas phase acetic acid will be removed from the atmosphere. Thus, based on these values, heterogeneous uptake of organic acids on mineral aerosol may be a significant, yet limited, removal mechanism for carboxylic acids in troposphere. Additionally, our results indicate that atmospheric processing of swelling clay minerals with water vapor and organic acids results in modification of the chemical composition of the gas phase and the entrained clay particle during transport through the atmosphere, which may lead to modified cloud properties such as the ability of these particles to become giant CCN or ice nuclei in addition to a possible modification of the optical properties of these minerals in the atmosphere.

Acknowledgements. This work was supported by the National Science Foundation under grant ATM – 0137261. CDH also received funding from the CIRES graduate student fellowship.

Edited by: J. Thornton

References

- Al-Abadleh, H. A. and Grassian, V. H.: Heterogeneous reaction of NO_2 on hexane soot: A Knudsen Cell and FT-IR Study, *J. Phys. Chem. A*, 104, 11 926–11 933, 2000.
- Al-Abadleh, H. A. and Grassian, V. H.: Oxide surfaces as environmental interfaces, *Surf. Sci. Rep.*, 52, 63–161, 2003.
- Al-Hosney, H. A., Carlos-Cuellar, S., Baltrusaitis, J., and Grassian, V. H.: Heterogeneous uptake and reactivity of formic acid on calcium carbonate particles: A Knudsen cell reactor, FT-IR and SEM study, *Phys. Chem. Chem. Phys.*, 7, 3587–3595, 2005.
- Anderson, D. M., Schwarz, M. J., and Tice, A. R.: Water-vapor adsorption by sodium montmorillonite at -5°C , *Icarus*, 34, 638–644, 1978.
- Bauer, S. E., Balkanski, Y., Schulz, M., Hauglustaine, D. A., and Dentener, F.: Global modeling of heterogeneous chemistry on mineral aerosol surfaces: Influence on tropospheric ozone chemistry and comparison to observations, *J. Geophys. Res.*, 109, D02304, doi:10.1029/2003JD003868, 2004.
- Bishop, J. L., Pieters, C. M., and Edwards, J. O.: Infrared spectroscopic analyses on the nature of water in montmorillonite, *Clays and Clay Min.*, 42, 702–716, 1994.

- Brunauer, S., Emmett, P. H., and Teller, E.: Adsorption of gases in multi-molecular layers, *J. Am. Chem. Soc.*, 60, 309–319, 1938.
- Carlos-Cuellar, S., Li, P., Christensen, A. P., Krueger, B. J., Burrichter, C., and Grassian, V. H.: Heterogeneous uptake kinetics of volatile organic compounds on oxide surfaces using a Knudsen cell reactor: Adsorption of acetic acid, formaldehyde, and methanol on α -Fe₂O₃, α -Al₂O₃, and SiO₂, *J. Phys. Chem. A*, 107, 4250–4261, 2003.
- Cases, J. M., Berend, I., Besson, G., Francois, M., Uriot, J. P., Thomas, F., and Poirier, J. E.: Mechanism of Adsorption and desorption of water-vapor by homoionic montmorillonite: 1. The sodium-exchanged form, *Langmuir*, 8, 2730–2739, 1992.
- DeBell, L. J., Vozzella, M., Talbot, R. W., and Dibb, J. E.: Asian dust storm events of spring 2001 and associated pollutants observed in New England by the Atmospheric Investigation, Regional Modeling, Analysis and Prediction (AIRMAP) monitoring network, *J. Geophys. Res.*, 109, D01304, doi:10.1029/2003JD003733, 2004.
- de Reus, M., Dentener, F., Thomas, A., Borrmann, S., Strom, J., and Lelieveld, J.: Airborne observations of dust aerosol over the North Atlantic Ocean during ACE 2: Indications for heterogeneous ozone destruction, *J. Geophys. Res.*, 105, 15 263–15 275, 2000.
- Dogan, A. U., Dogan, M., Onal, M., Sarikaya, Y., Aburub, A., and Wurster, D. E.: Baseline studies of the Clay Minerals Society source clays: Specific surface area by the Brunauer Emmett Teller (BET) method, *Clays and Clay Min.*, 54, 62–66, 2006.
- Downing, H. D. and Williams, D.: Optical constants of water in the infrared, *J. Geophys. Res.*, 80, 1656–1661, 1975.
- Falkovich, A. H., Schkolnik, G., Ganor, E., and Rudich, Y.: Adsorption of organic compounds pertinent to urban environments onto mineral dust particles, *J. Geophys. Res.*, 109, D02208, doi:10.1029/2003JD003919, 2004.
- Flatau, P. J., Walko, R. L., and Cotton, W. R.: Polynomial fits to saturation vapor-pressure, *J. Appl. Meteorol.*, 31, 1507–1513, 1992.
- Frinak, E. K., Mashburn, C. D., Tolbert, M. A., and Toon, O. B.: Infrared characterization of water uptake by low temperature Namontmorillonite: Implications for Earth and Mars, *J. Geophys. Res.*, 110, D09308, doi:10.1029/2004JD005647, 2005.
- Frinak, E. K., Wermeille, S. J., Mashburn, C. D., Tolbert, M. A., and Pursell, C. J.: Heterogeneous reaction of gaseous nitric acid on gamma-phase iron (III) oxide, *J. Phys. Chem. A*, 108, 1560–1566, 2004.
- Grassian, V. H.: Heterogeneous uptake and reaction of nitrogen oxides and volatile organic compounds on the surface of atmospheric particles including oxides, carbonates, soot and mineral dust: Implications for the chemical balance of the troposphere, *Int. Rev. Phys. Chem.*, 20, 467–548, 2001.
- Haderlein, S. B. and Schwarzenbach, R. P.: Adsorption of substituted nitrobenzenes and nitrophenols to mineral surfaces, *Environ. Sci. Technol.*, 27, 316–326, 1993.
- Hall, P. L. and Astill, D. M.: Adsorption of water by homoionic exchange forms of Wyoming montmorillonite (SWy-1), *Clays and Clay Min.*, 37, 355–363, 1989.
- Hendricks, S. B., Nelson, R. A. and Alexander, L. T.: Hydration mechanism of the clay mineral montmorillonite saturated with various cations, *J. Am. Chem. Soc.*, 62, 1457–1464, 1940.
- Hensen, E. J. M. and Smit, B.: Why clays swell, *J. Phys. Chem. B*, 106, 12 664–12 667, 2002.
- Khan, I., Brimblecombe, P., and Clegg, S. L.: Solubilities of pyruvic acid and the lower (C₁–C₆) carboxylic acids: Experimental determination of equilibrium vapor pressures above pure aqueous solutions, *J. Atmos. Chem.*, 22, 285–302, 1995.
- Kubicki, J. D., Schroeter, L. M., Itoh, M. J., Nguyen, B. N., and Apitz, S. E.: Attenuated total reflectance Fourier-transform infrared spectroscopy of carboxylic acids adsorbed onto mineral surfaces, *Geo. Cosmo. Acta*, 63, 2709–2725, 1999.
- Lee, S. H., Murphy, D. M., Thomson, D. S., and Middlebrook, A. M.: Chemical components of single particles measured with Particle Analysis by Laser Mass Spectrometry (PALMS) during the Atlanta SuperSite Project: Focus on organic/sulfate, lead, soot and mineral particles, *J. Geophys. Res.*, 107, 4003, doi:10.1029/2000JD000011, 2002.
- Lee, S. H., Murphy, D. M., Thomson, D. S., and Middlebrook, A. M.: Nitrate and oxidized organic ions in single particle mass spectra during the 1999 Atlanta Supersite Project, *J. Geophys. Res.*, 108, 8417, doi:10.1029/2001JD001455, 2003.
- Li, P., Perreau, K. A., Covington, E., Song, C. H., Carmichael, G. R., and Grassian, V. H.: Heterogeneous reactions of volatile organic compounds on oxide particles of the most abundant crustal elements: Surface reactions of acetaldehyde, acetone, and propionaldehyde on SiO₂, Al₂O₃, Fe₂O₃, TiO₂, and CaO, *J. Geophys. Res.*, 106, 5517–5529, doi:10.1029/2000JD900573, 2001.
- Lide, D. R. (Ed.): *Handbook of Chemistry and Physics*, CRC Press, Boca Raton, FL, 2004.
- Madejova, J. and Komadel, P.: Baseline studies of The Clay Minerals Society Source Clays: Infrared methods, *Clays and Clay Min.*, 49, 410–432, 2001.
- Marti, J. and Mauersberger, K.: A survey and new measurements of ice vapor pressure at temperatures between 170 and 250 K, *Geophys. Res. Lett.*, 20, 363–366, 1993.
- Mashburn, C. D., Frinak, E. K., and Tolbert, M. A.: Heterogeneous uptake of nitric acid on Na-montmorillonite clay as a function of relative humidity, *J. Geophys. Res.*, 111, D15213, doi:10.1029/2005JD006525, 2006.
- Max, J. J. and Chapados, C.: Infrared spectroscopy of aqueous carboxylic acids: Comparison between different acids and their salts, *J. Phys. Chem. A*, 108, 3324–3337, 2004.
- Maxwell-Meier, K., Weber, R., Song, C., Orsini, D., and Ma, Y.: Inorganic composition of fine particles in mixed mineral dust-pollution plumes observed from airborne measurements during ACE-Asia, *J. Geophys. Res.*, 109, D19S07, doi:10.1029/2003JD004464, 2004.
- McKendry, I. G., Hacker, J. P., and Stull, R.: Long-range transport of Asian dust to the Lower Fraser Valley, British Columbia, Canada, *J. Geophys. Res.*, 106(D16), 18 361–18 370, 2001.
- Mooney, R. W., Keenen, A. G., and Wood, L. A.: Adsorption of water vapor by montmorillonite: I. Heat of desorption and application of BET Theory, *J. Am. Chem. Soc.*, 74, 1367–1371, 1952.
- Newman, A. C. D.: The specific surface area of soils determined by water sorption, *Soil Sci.*, 34, 23–32, 1983.
- Pennell, K. D., Rhue, R. D., Rao, P. S. C., and Johnston, C. T.: Vapor phase sorption of para-xylene and water on soils and clay minerals, *Environ. Sci. Technol.*, 26, 756–763, 1992.
- Ravishankara, A. R.: Heterogeneous and multiphase chemistry in the troposphere, *Science*, 276, 1058–1065, 1997.
- Reid, E. A., Reid, J. S., Meier, M. M., Dunlap, M. R., Cliff, S.

- S., Broumas, A., Perry, K., and Maring, H.: Characterization of African dust transported to Puerto Rico by individual particle and size segregated bulk analysis, *J. Geophys. Res.*, 108, 8591, doi:10.1029/2002JD002935, 2003.
- Russell, L. M., Maria, S. F., and Myneni, S. C. B.: Mapping organic coatings on atmospheric particles, *Geophys. Res. Lett.*, 29(16), 1779, doi:10.1029/2002GL014874, 2002.
- Shilling, J. E., Connelly, B. M., and Tolbert, M. A.: Uptake of small oxygenated organic molecules onto ammonium nitrate under upper tropospheric conditions, *J. Phys. Chem. A*, 110, 6687–6695, 2006.
- Shilling, J. E. and Tolbert, M. A.: Uptake of acetic acid on thin ammonium nitrate films as a function of temperature and relative humidity, *J. Phys. Chem. A*, 108, 11 314–11 320, 2004.
- Sokolik, I. N., Toon, O. B., and Bergstrom, R. W.: Modeling the radiative characteristics of airborne mineral aerosols at infrared wavelengths, *J. Geophys. Res.*, 103, 8813–8826, 1998.
- Sun, Y., Zhuang, G., Wang, Y., Zhao, X., Li, J., Wang, Z., and An, Z.: Chemical composition of dust storms in Beijing and implications for the mixing of mineral aerosol with pollution aerosol on the pathway, *J. Geophys. Res.*, 110, D24209, doi:10.1029/2005JD006054, 2005.
- Talbot, R. W., Mosher, B. W., Heikes, B. G., Jacob, D. J., Munger, J. W., Daube, B. C., Keene, W. C., Maben, J. R., and Artz, R. S.: Carboxylic acids in the rural continental atmosphere over the eastern United States during the Shenandoah Cloud and Photochemistry Experiment, *J. Geophys. Res.*, 100(D5), 9335–9344, doi:10.1029/95JD00507, 1995.
- Tang, Y. H., Carmichael, G. R., Kurata, G., Uno, I., Weber, R. J., Song, C. H., Guttikunda, S. K., Woo, J. H., Streets, D. G., Wei, C., Clarke, A. D., Huebert, B., and Anderson, T. L.: Impacts of dust on regional tropospheric chemistry during the ACE-Asia experiment: A model study with observations, *J. Geophys. Res.*, 109, D19S21, doi:10.1029/2003JD003806, 2004.
- Underwood, G. M., Li, P., Usher, C. R., and Grassian, V. H.: Determining accurate kinetic parameters of potentially important heterogeneous atmospheric reactions on solid surfaces with a Knudsen Cell reactor, *J. Phys. Chem. A*, 104, 819–829, 2000.
- Van Olphen, H. and Fripiat, J. J.: Data handbook for clay minerals and other non-metallic minerals, Pergamon Press, Oxford, England, 1979.
- Wang, Y., Zhuang, G., Sun, Y., and An, Z.: Water-soluble part of the aerosol in the dust storm season: Evidence of the mixing between mineral and pollution aerosols, *Atmos. Environ.*, 39, 7020–7029, 2005.
- Zhang, X. Y., Arimoto, R., and An, Z. S.: Dust emission from Chinese desert sources linked to variations in atmospheric circulation, *J. Geophys. Res.*, 102, 28 041–28 047, 1997.
- Zhang, Y. and Carmichael, G. R.: The role of mineral aerosol in tropospheric chemistry in East Asia – A model study, *J. Appl. Meteorol.*, 38, 353–366, 1999.